

MAP NO. 2 ADULT CHINOOK AND COHO MIGRATION ROUTES

Seasonality Information - Adult chinook and coho salmon and steelhead.


SPECIES	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Chinook	Some always present in inshore waters.				Adults return to spawning streams.							
Coho	Some always present in inshore waters.				Adults return to spawning streams.							
Steelhead	Adults present in inshore waters all year round											

References


- McAllister, C. pers. com. DFO - Pacific Biological Station, Nanaimo, B.C.
- Milne, R.J. 1957. Recent B.C. Spring Salmon and Coho Salmon Tagging Experiments and a Comparison 1925-1930. Fish. Res. Board Can. Bull. 113: 56 p.

Explanatory Notes

Migration routes of adult chinook and coho are determined through the analysis of commercial fisheries catch statistics and through tagging studies. Adult fish are caught in seines or by trolling, marked with coded tags and released. The movements of the tagged fish from the tagging sites can be interpolated when they are recaptured in subsequent surveys or in the commercial and recreational fishery.



- LEGEND**
- Chinook Migration Routes
 - Coho Migration Routes
 - Chinook Tagging Sites
 - Coho Tagging Sites



Department of Fisheries and Oceans, Pacific Region,
Habitat Management Division
February, 1985

