Morice Watershed-based Fish Sustainability Planning

A Plan to Conserve and Protect Morice Watershed Fish Populations and their Habitat – Stage II

Greg Tamblyn¹ and Chad Croft²

September 2003

Prepared for Fisheries and Oceans Canada &
Community Futures Development Corporation of Nadina

Funded by: Fisheries and Oceans Canada - Habitat Conservation and Stewardship Program





^{1 -} Independent Consultant

² – Gartner Lee Ltd.

ACKNOWLEDGEMENTS

The WFSP process, like other strategic planning processes, requires motivation, dedication and organization. As such, the authors would like to thank the members of the Morice WFSP planning and technical committees for their time and efforts over the past 18 months. Their guidance throughout, combined with their comments on draft materials have been integral to completing the plan to this stage. We also appreciate the efforts of Dale Gueret and Bruce Shepherd, who provided valuable comments on the draft plan.

We would especially like to thank Martin Forbes of Fisheries and Oceans Canada for seeing the benefits of this planning process from the beginning and for his leadership in funding this project. Funding was provided by the Habitat Conservation and Stewardship Program of Fisheries and Oceans Canada. Fisheries Renewal BC provided seed funding during the initial year of the project.

TABLE OF CONTENTS

ACKNOWLEDGEMENTSI
TABLE OF CONTENTSII
LIST OF TABLESV
LIST OF FIGURESV
PREFACEVI
1.0 INTRODUCTION1
1.1 What is Watershed-based Fish Sustainability Planning?
1.2 Rationale for adopting a WFSP approach
1.3 The Morice Watershed-based Fish Sustainability Plan 2 1.3.1 Guiding Principles and Process Goals 2 1.3.2 Stage I – The Bulkley WFSP 3 1.3.3 Stage II – The Morice WFSP 3 1.3.4 Public Participation 6 1.3.5 Strategic Direction 6 1.3.6 Time frame for the Morice WFSP 6 1.3.7 Linkages to other Planning Processes 7
2.0 WATERSHED PROFILE9
2.1 Physiography
2.2 Hydrology
2.3 Water Quality
2.4 Land and Water Use 10 2.4.1 Forest Management 10 2.4.2 Mineral Resource Development 10 2.4.3 Transportation and Utilities 10 2.4.4 Other Land Use 11 2.4.5 Water Use 11 2.4.6 Settlements 11 2.4.7 Employment and Income 11
3.0 FISHERIES RESOURCES12
3.1 Sockeye Salmon

Morice Watershed-based Fish Sustainability Plan - Stage II

3.2 Chinook Salmon	14
3.2.1 Population Status and Trends	14
3.2.2 Limiting Factors to Production	14
3.2.3 Habitat Condition and Protection	14
3.3 Coho Salmon	15
3.3.1 Population Status and Trends	13
3.3.2 Limiting Factors of Production	15
3.3.3 Habitat Condition and Protection	15
	1.
3.4 Pink Salmon	10
3.4.1 Population Status and Trends	10
3.4.2 Limiting Factors on Production	
3.4.3 Habitat Condition and Protection	10
3.5 Steelhead	1
3.5.1 Population Status and Trends	
3.5.2 Limiting Factors for Production	
3.5.3 Habitat Condition and Protection	
J.J.J Habitat Condition and Protection	
3.6 Rainbow Trout	18
3.6.1 Population Status and Trends	18
3.6.2 Limiting Factors of Production	18
3.6.3 Habitat Condition and Protection	18
3.7 Cutthroat Trout	18
3.7.1 Population Status and Trends	18
3.7.2 Habitat Condition and Protection	18
3.8 Bull Trout	10
3.8.1 Population Status and Trends	19
3.8.2 Limiting Factors for Production	19
3.8.3 Habitat Condition and Protection	19
J.0.5 1140144 Condition 414 11000101	
3.9 Dolly Varden	20
3.9.1 Population Status and Trends	20
3.9.2 Habitat Condition and Protection	20
3.10 Lake Trout	
3.11 Kokanee	
5.11 Nokanee	
3.12 Whitefish	20
3.13 Other species	20
3.13.1 Burbot	20
3.13.2 Lake Chub	21
3.13.3 Longnose Sucker	
3.13.4 Largescale Suckers	
3.13.5 Redside Shiners	
3.13.6 Longnose Dace	21
3.13.7 Prickly Sculpin	21
3.13.8 Pacific Lamprey	
4.0 FACTORS AFFECTING MORICE FISH POPULATIONS	22

Morice Watershed-based Fish Sustainability Plan - Stage II

4.1 Commercial and In-river Fishing	22
4.2 Land Use Activities	22
4.3 Fish Farms	22
5.0 PRIORITY ISSUES	23
5.1 Morice Watershed Issues	2 3
5.2 Selection of Priority Issues	24
6.0 OBJECTIVES AND STRATEGIES FOR SELECTED PRIORITY ISSUES	25
7.0 MONITORING FRAMEWORK	30
7.1 What is Monitoring?	30
7.2 Why is Monitoring Important?	30
7.3 Monitoring for the Morice WFSP	30
7.4 The Monitoring Toolbox	33
7.5 Data Management	34
8.0 IMPLEMENTATION FRAMEWORK	35
9.0 NEXT STEPS	35
10.0 REFERENCES CITED	37
APPENDIX A Members of the Morice WFSP planning & technical committee	s39
APPENDIX B Land use analysis table	40
APPENDIX C1 General issues table	41
APPENDIX C2 Species Specific Issues	42
APPENDIX D Possible scoring matrix to rank issues	47

LIST OF TABLES

Table 1.	Basic sector employment estimates as a percentage of total jobs for the Morice Timber Supply Area (including Indian Reserves), 1996
Table 2.	Number of people employed in Houston and average earnings based on 2001 statistics.
Table 3.	Priority fish species and stocks of interest
Table 4.	Possible tools for monitoring forest development (adapted from Mackay 1998) 33
LIST C	OF FIGURES
Figure 1:	The sub-planning units of the Bulkley watershed for Stage I of the Bulkley WFSP 4
	The Morice watershed.
	Links among land use planning processes related to the Morice WFSP
Figure 4:	Relative impacts of land use within the Morice River watershed
Figure 5:	Possible steps to develop a monitoring plan. Source Mackay (1998) with minor modifications

PREFACE

The Morice Watershed-based Fish Sustainability Plan is a work in progress. This report is the culmination of efforts over 18 months toward developing a plan to conserve and protect the fish and fish habitat of the Morice River watershed. This document will act as a starting point when watershed planning resumes for the area. Significant work is required to complete the process as described in the Watershed-based Fish Sustainability Planning Guidebook (MELP & DFO 2001) and to develop a thorough and effective plan.

The Ministry of Sustainable Resource Management is currently funding a review of Watershed-based Fish Sustainability Planning (WFSP). The purpose of the review is to clarify the nature of federal and provincial government involvement in WFSP and to determine how to implement the process effectively throughout the province given recent widespread reorganization and new policy direction.

1.0 INTRODUCTION

1.1 What is Watershed-based Fish Sustainability Planning?

Watershed-based Fish Sustainability Planning (WFSP) is a planning process focused on fish populations and aquatic habitat. "Its overall goal is to ensure effective long-term conservation of fish and fish habitat" (MELP & DFO 2001). The process, introduced in 2001, is described in a guidebook developed by provincial and federal government agencies responsible for managing fish in British Columbia. This guidebook outlines a four stage planning sequence that can be used by interested First Nations, community groups, industries and government agencies to identify and protect those fish populations and fish habitats that most urgently require attention:

- Stage I: Establishes regional priorities for conservation of significant fish populations and habitats based on a broad profile of a large watershed or sub-basin. The stage concludes with the selection of one or more candidate watershed planning units for more detailed planning.
- Stage II: Establishes priorities and an action plan framework for key planning units identified in Stage I based on a detailed watershed profile and social, economic and political considerations. In this stage, strategic direction (goals), management objectives and strategies are established.
- Stage III: Develops an action plan detailing how the plan will be implemented and monitored, and by whom.
- Stage IV: Implements, monitors and modifies the plan.

1.2 Rationale for adopting a WFSP approach

Watershed-based Fish Sustainability Planning was adopted for the Bulkley watershed with the goal of completing strategic watershed planning begun by the Bulkley-Morice Salmonid Preservation Group (BMSPG). The BMSPG, formed as the delivery group for Fisheries Renewal BC in the Bulkley watershed, consisted of representatives from First Nations and community organizations concerned about fish stocks and the health of the watershed. The group was chaired and administered by Community Futures Development Corporation of Nadina (Nadina Community Futures).

The BMSPG completed its first stage of watershed planning in mid-2001. The resulting document entitled *Healthy Watersheds, Healthy Communities: Bulkley-Morice Salmonid Preservation Group Draft Strategic Plan – Phase 1* (Tamblyn and Donas 2001) identified eight priority watershed issues and listed draft goals, objectives and strategies to address each issue. To maximize the plan's effectiveness as a direction-setting document, further planning was required to determine priority sub-watersheds, and to develop an implementation and monitoring strategy. The BMSPG felt that integrating the existing draft plan into a government endorsed WFSP process would achieve these goals within guidelines designed to create consistency among watershed plans throughout the province.

1.3 The Morice Watershed-based Fish Sustainability Plan

1.3.1 Guiding Principles and Process Goals

Nine guiding principles and several process goals provide the framework for the Morice WFSP:

- a) The Project will be undertaken with a focus on the sustainability of wild fish stocks, and with attention to both anadromous and non-anadromous species, genetic diversity, and commercial and non-commercial species;
- b) The Project will maintain a focus on watersheds, including their processes and their interconnections, both instream and upland;
- c) The Project will be carried out using a "fish first" approach; that is, with an emphasis on the needs of fish;
- d) The Project will identify priorities for the protection and restoration of fish stocks and habitat;
- e) The Project will build on existing initiatives;
- f) The Project will use the best information currently available, including scientific data, traditional ecological knowledge, land and resource development trends, and community values;
- g) The Project will identify data gaps and will provide recommendations on priorities and means of filling those gaps;
- h) The Project will use "adaptive management"; that is, an approach that incorporates ongoing monitoring and assessment to create a living plan that has no defined completion point; and
- i) The Project will be carried out using an inclusive and consensus-based process that openly involves members of the community, including residents, responsible agencies, and commercial and non-commercial users of the watershed.

Additional process goals are:

- To integrate WFSP with concurrent land use planning processes, including the Morice Land and Resource Management Plan (LRMP) and the Morice-Lakes Innovative Forest Practices Agreement (IFPA), to ensure coordination and to minimize overlap;
- To articulate the connection between managing for sustainable fish populations and managing for sustainable land use; and
- To develop a modular plan for the benefit of implementing industry and agencies.

1.3.2 Stage I - The Bulkley WFSP

Nadina Community Futures and Fisheries and Oceans Canada initiated WFSP for the Bulkley watershed soon after the completion of Phase I of the BMSPG plan. Stage 1 of the WFSP was launched in late 2001 with the establishment of a planning committee (see Appendix A). As an initial step, the planning committee divided the Bulkley watershed into four planning units: the Morice watershed, and the upper, middle and lower Bulkley watersheds (Figure 1). Following consultation with the public, First Nations, and government via questionnaires, the committee selected the Morice River watershed as the priority planning unit for continued planning (Figure 2).

Around the same time, an independent Skeena stage I WFSP process, led by the Skeena Fisheries Commission, selected the Morice watershed as one of the three most important watersheds in the Skeena from a salmon production perspective (Gottesfeld et al. 2002).

1.3.3 Stage II - The Morice WFSP

Stage II began in January 2002 with the formation of the Morice WFSP technical committee (see Appendix A). The planning and technical committees mapped out Stage II of the planning process with the help of the WFSP guidebook. The technical committee's first task was to guide the creation of the Morice biophysical watershed profile (Bustard and Schell 2002). This document summarized the fish stocks in the watershed including life histories, population status and trends, distributions and key habitats, probable limiting factors to fish productivity, and information gaps. To aid the WFSP process further, Croft and Bahr (2002) developed a problem analysis to attempt to link fish and fish habitat trends with resource management practices. In addition, Croft conducted a GIS-based land use analysis using Watersheds BC data¹ to provide a baseline for future analysis and monitoring (see Section 4.2 and Appendix B).

The technical committee then categorized issues and concerns identified by the public and Bustard and Schell (2002). General aquatic ecosystem / watershed issues are summarized in Appendix C1. Species-specific issues are listed in Appendix C2. When the Morice WFSP process continues, the criteria suggested in Appendix D may help rank the issues into a priority list.

For this current draft of the plan, key issues were extracted from the conclusions and recommendation sections of Bustard and Schell (2002). Draft objectives, example strategies and possible indicators were identified for selected priority issues (Section 6).

¹ A detailed land use analysis expected from the IFPA did not materialize prior to drafting this report.




Figure 1: The sub-planning units of the Bulkley Watershed for Stage I of the Bulkley WFSP.






Figure 2: The Morice watershed.

1.3.4 Public Participation

Burnout associated with multiple ongoing planning processes, combined with increasing demands on volunteers, resulted in limited direct public participation at planning and technical committee meetings. Nonetheless, we made attempts to ensure the incorporation of public concerns into the Morice WFSP through:

- Regular review of the BMSPG draft strategic plan (which was a public process);
- Review of compiled lists of issues and concerns from the Morice LRMP Fish and Fish Habitat Sector;
- Presence of one of the authors at Morice LRMP Fish and Fish Habitat Sector meetings;
 and
- A presentation on WFSP and distribution of meeting summaries to Morice LRMP Fish and Fish Habitat Sector members.

1.3.5 Strategic Direction

The aquatic ecosystems and most fish populations in the Morice River watershed appear to be relatively healthy. Unlike many watersheds identified as priorities for watershed planning in British Columbia, tremendous opportunities exist to protect existing habitats and fish stocks in the Morice. While rehabilitating damaged habitat is required in some areas, proactive, preventative actions are preferred where possible. The overall strategic directions for the Morice WFSP are:

- To maintain natural aquatic ecosystem functions and processes; and
- To maintain or increase populations of endemic wild fish populations to levels equal to the productive capacity of the Morice watershed.

More detailed goals are:

Biological goals:

- To protect existing habitat and rehabilitate impacted habitat where identified as a priority;
- To ensure that aquatic species do not decline below sustainable levels (limit reference points where established); and
- To allow adequate escapements of anadromous fish to fully seed the available habitat, while maintaining a natural species mix and balance.

Socio-economic goals:

- To maintain or improve quality of life for communities in the Bulkley and Skeena watersheds through ensuring clean water supplies and providing world class recreational angling opportunities; and
- To provide long-term economic opportunities for commercial and First Nation fishers and the local tourism industry by maintaining or increasing fish populations and fish habitat.

1.3.6 Time frame for the Morice WFSP

We expect the completion of Stage II and Stage III of the Morice WFSP to take another year. Implementation will require several additional years, depending on opportunities to implement

some of the plan's objectives through the Morice LRMP, the Morice-Lakes Innovative Forest Practices Agreement and Sustainable Resource Management Plans.

The Morice WFSP is a living plan meant to be altered based on implementation and effectiveness monitoring and society's changing priorities. We envision significant updating and revision to occur every 5-10 years.

1.3.7 Linkages to other Planning Processes

The Morice WFSP is facing unique challenges and opportunities resulting from concurrent planning processes in the area. The ongoing Morice Land and Resource Management Plan and the Morice-Lakes Innovative Forest Practices Agreement (IFPA) both encompass the WFSP planning area. Stage I of the Skeena WFSP process also incorporated the Morice watershed. With so many competing processes, the main challenges lie in 1) avoiding duplication and in 2) attracting participants from industry, government and the general public to the WFSP. Nevertheless, opportunities abound for synergism due to the niches of each planning process. Figure 3 outlines the connections among ongoing related planning processes.

WFSP and LRMP

The fish sector of the Morice LRMP is providing input into the Morice WFSP. In return, the fish sector and the Government Technical Team use technical products from the WFSP for the LRMP. Should the Morice WFSP be completed promptly, we hope that some of the strategic goals and objectives for the WFSP will be incorporated into the Morice LRMP.

WFSP and IFPA

Members of the IFPA and WFSP are working collaboratively to the benefit of both processes. The IFPA and subsequent Sustainable Forest Management Plans may provide avenues for monitoring and implementing part of the WFSP. As the Morice WFSP continues, the technical committee can utilize modelling and land use analyses conducted by the IFPA, while the WFSP can help fill aquatic management and monitoring gaps in the IFPA. Together, these processes will link sustainable fish habitat and populations to sustainable land use management.

Morice and Skeena WFSP

The Morice WFSP planning committee has endeavored to work collaboratively with other WFSPs in the region, including the Skeena Stage I process. The other two recommended Stage II processes in the Skeena watershed - the Kispiox and Lakelse - have been postponed due to funding shortages. However, in the future, we hope to achieve some consistency among plans and streamline requests for help and information from government agencies by working together.




Figure 3: Links among land use planning processes related to the Morice WFSP.

2.0 WATERSHED PROFILE

2.1 Physiography

The Morice watershed, located southwest of Houston, B.C. (Figure 2), is the largest tributary to the Bulkley River, with a catchment area of 4,349 km². This basin drains both the interior plateau south of Houston, and the Coast Mountains to the west. The Morice River, headed by Morice Lake, flows northeast for approximately 80 km before entering the Bulkley River four kilometers west of Houston. The Bulkley then proceeds approximately 150 km northwest to join the Skeena River at Hazelton.

The predominant biogeoclimatic zone covering most of the lowland coniferous forests in the watershed is the Sub-Boreal Spruce (SBS) zone with dry cool (dk), moist cold (mc), and wet cool (wk) sub-zones. The SBS zone meets the Englemann Spruce-Subalpine Fir (ESSF) zone at upper elevations ranging from 900 to 1300 m. The ESSF is dominated by continuous forests stretching into sub alpine meadows at upper elevations and is characterised by moist cold (mc), moist cool (mk) and moist, very cold (mv) sub-zones. Additional biogeoclimatic zones and sub-zones include the Alpine Tundra (AT) at the highest elevations, Coastal Western Hemlock, wet submaritime (CWHws), flanking the shores of the large lakes in the southwestern portion of the watershed, and Mountain Hemlock, moist maritime (MHmm), in the southern-most headwaters.

2.2 Hydrology

The Morice River hydrograph is primarily influenced by snowmelt, with peak flows occurring from late May through July. Historically, June experiences the highest average monthly flows, which can approach 500 m³/second. In some years, rain on snow events produce peak flows in October and early November (Bustard and Schell 2002).

Two of the largest tributaries within the Morice River watershed are the Nanika (895 km²) and Atna (300 km²) rivers, which empty into Morice Lake. The Thautil River and Gosnell Creek combine to form the largest tributary (535 km²) downstream of Morice Lake. Hydrologically, this is one of the most important systems in the Morice watershed as it influences peak flows and contributes significant bedload and sediment to the Morice River mainstem (Bustard and Schell 2002).

Due to slow glacier melt, abundant lake storage and fall rains, the mainstem Morice and Nanika rivers, as well as the larger tributaries such as the Thautil, Gosnell and Houston Tommy, maintain relatively high flows throughout the year until freeze up, typically in November. The lowest stream flows in both the mainstem rivers and smaller tributaries occur during the late winter usually from March to the middle of April at the higher elevations. Smaller, lower elevation streams draining the Interior Plateau, such as McBride, Lamprey, and Owen creeks often experience low summer flows (Bustard and Schell 2002).

2.3 Water Quality

Although the overall nutrient level is low, Morice River water quality is considered excellent, with a typically clear water column, a near neutral pH level, a mean alkalinity of 23 mg/l and a mean conductivity of 53 μ ohms/cm (Remington 1996; Gottesfeld et al. 2002; Bustard and Schell 2002).

Impacts to water quality in the Morice have been linked to both forestry and mining land use activities. Bustard (1996) considered increased suspended sediments associated with road construction and inadequate road maintenance to be the greatest logging related impact on Morice watershed streams. Remington (1996) identified the Silver Queen Mine on the eastern shore of Owen Lake as a source of elevated zinc (Zn) and copper (Cu) levels in the Owen watershed.

2.4 Land and Water Use

Industrial land use in the Morice watershed is dominated by forest management activities, an expansive road network, and past mineral exploration. Other land use activities include recreation, tourism, trapping, guide outfitting, cattle grazing and utility corridors.

2.4.1 Forest Management

Forest management activities in the Morice watershed have historically involved progressive road development and well distributed timber harvesting, typically characterized by clearcut silviculture systems.

The current mountain pine beetle infestation in the Morice has resulted in a shift in forest management strategies. Recent forest development activities have focused on small scale harvesting of infested stands with a greater reliance on temporary roads. However, should the infestation reach epidemic proportions, extensive harvesting control strategies present a risk to the integrity of the aquatic ecosystems of the watershed.

2.4.2 Mineral Resource Development

The mining industry has a long history in the area, as evidenced by extensive mineral exploration. However, of five developed prospects, only one mine has operated within the watershed. The Silver Queen Mine, located on the eastern shore of Owen Lake, produced silver, gold, cadmium, copper, lead, and zinc in 1972 and 1973 (Horn and Tamblyn 2000).

2.4.3 Transportation and Utilities

Progressive construction by the forest industry has resulted in a network of roads throughout most of the valley bottoms and onto the majority of the upland plateaus. This road network is primarily utilized by the forest industry, recreationalists, trappers, guide-outfitters and the mining industry. Since 1996, Huckleberry Mine, located 120 km southwest of Houston, has used the Morice-Owen Road to haul ore. A BC hydro line servicing Huckleberry Mine closely follows the road right of way (Gottesfeld et al. 2002).

(

2.4.4 Other Land Use

The Morice watershed is used extensively for recreational activities such as hunting, fishing, hiking, camping, canoeing and snowmobiling. Small capital ventures such as guide outfitting, trapping and tourism also utilize the watershed. Cattle range tenures exist near Houston, to the east of Owen Lake, and to the east of Tagetochlain Lake (Horn and Tamblyn 2000).

2.4.5 Water Use

Water use in the Morice watershed is restricted to 14 water licences that authorize the use of water from Morice River tributaries for domestic, stock watering, waterworks and enterprise as defined by Land and Water BC. The total allowable consumption authorized under the fourteen water licences is 35,200 gallons per day (GD) with 7000 GD for domestic, 10,700 GD for stock watering, 10,500 GD for enterprise and 7,000 GD for waterworks (Land and Water BC Inc. 2003).

2.4.6 Settlements

The entire Morice watershed is located within the Wet'suwet'en traditional territory. Wet'suwet'en people resided at various village sites throughout the watershed until the early 1950's. Currently, less than 20 people reside permanently within the watershed (Gottesfeld et al. 2002).

The District of Houston, located four kilometres east of the Morice and Bulkley River confluence, is the primary settlement and is home to approximately 3600 people (2001 statistics). Houston is experiencing a decline in population and lost 9.1% of its population between 1996 and 2001 (Stats. Can 2003). Neighbouring communities utilizing the Morice watershed for work or recreation include Smithers, Telkwa, and Topley.

2.4.7 Employment and Income

The District of Houston and neighbouring communities rely heavily on natural resources for employment. In 1996, half the jobs in the Morice Timber Supply Area (includes Houston, Granisle and Topley) were with the forest industry (Table 1). Other important employers were the public sector, tourism, construction, agriculture and mining. These statistics are aging and under estimate current numbers for the mining industry, and over estimate the size of the public sector. Significant cutbacks in 2002 and 2003 drastically reduced the provincial government work force. Average wages for employees in the District of Houston are summarized in Table 2.

Table 1. Basic sector employment estimates as a percentage of total jobs for the Morice Timber Supply Area (including Indian Reserves), 1996.

Sector	% Basic Employment
Forestry	50%
Mining ²	3%
Public sector	22%
Tourism	9%
Agriculture	6%
Construction	7%
Other ^b	2.1%

- a. Note, the percentages shown above reflect each sector's share of an area's total employment, including direct and indirect employment.
- b. The "other" category includes several sectors, including transportation, fishing, trapping, and high technology
- c. Basic income flows into the community from outside of the area. Non-basic employment refers to local income that occurs due to the spending of basic income in the local area (e.g., local goods and services)

(Source: Horn and Tamblyn 2000)

Table 2. Number of people employed in Houston and average earnings based on 2001 statistics.

Characteristics	Numbers and earnings
# persons with earnings	2090
Average earnings	\$34,338
# persons working full time	980
Average earnings for full time workers	\$48,862

(Stats. Can. 2003).

3.0 FISHERIES RESOURCES

The Morice watershed is well known for its high fisheries values and is a major producer of all Pacific salmon species except chum (*Oncorhynchus keta*). The watershed forms an integral part of the salmon production in the Skeena drainage, and is therefore important to First Nations groups, as well as commercial and recreational fisheries. The Morice is also famous for its summer steelhead (*O. mykiss*) run.

In addition to producing Pacific salmon, the Morice watershed is utilized by rainbow trout (O. mykiss), cutthroat trout (O. clarki), kokanee (O. nerka), bull trout (Salvelinus confluentus), Dolly Varden (S. malma) and lake trout (S. namaycush). Numerous other resident fish species including mountain whitefish (Prosopium williamsoni), burbot (Lota lota), lake (Conesius plumbeus) and peamouth (Mylocheilus caurinus) chub, longnose (Catostomus catostomus) and

² The figures for basic mining employment and after-tax income related to mining do not include the Huckleberry mine, which opened in 1997, a year after these statistics were derived.

largescale (C. macrocheilus) suckers, redside shiners (Richardsonius balteatus), longnose dace (Rhinichthys cataractae), prickly sculpins (Cottus asper) and pacific lamprey (Lampetra tridentata) can also be found in the watershed.

3.1 Sockeye Salmon

3.1.1 Population Status and Trends

Morice-Nanika sockeye is the only significant sockeye run in the Morice watershed. As part of the Skeena watershed escapement, the Morice-Nanika sockeye stock comprises 1-2 % on average, while the Babine stock makes up 91%. The overwhelming difference in size of the two stocks, in conjunction with commercial fishing methods and management, makes the Morice-Nanika stock vulnerable to over harvesting by commercial fishing operations targeting the larger Babine stock (Bustard and Schell 2002).

The Morice-Nanika sockeye run has generally been in decline since the mid-1950's. Returns in the early 1950's averaged 50,000 fish; however, by 1955 escapement levels dropped to 4,000, and remained less than 6,000 fish until the early 1990's. From 1991 to 1997 the stock appeared to rebound with returns of 22,000 – 41,000 fish; however, by 1998, numbers dropped again and remained low through to 2001 (Bustard and Schell 2002).

The drop in numbers is a particular concern for the Office of the Wet'suwet'en and the Moricetown Band who rely on Nanika sockeye at Moricetown Canyon for a First Nation's food fishery.

3.1.2 Limiting Factors to Production

The factors influencing production and possibly contributing to the sharp decline and continued low abundance of Morice-Nanika sockeye since the 1950's are thought to be the low nutrient levels within Morice Lake, low spawner recruitment, climatic conditions causing flooding, redd dewatering and freezing, predation and possibly the availability of quality spawning habitat during high escapement years (Bustard and Schell 2002).

3.1.3 Habitat Condition and Protection

Critical sockeye habitat in the Morice is limited to a few key locations in the upper portion of the watershed. The only significant spawning area for Morice-Nanika Sockeye is located in the upper Nanika River. This area is relatively deep and subject to stable warm water flows from Kidprice Lake. Egg survival is expected to be good in these spawning sites due to a lack of ice, flooding and sediment inputs. Secondary spawning locations downstream of Glacier Creek may be subject to changing hydrological flows, increase sediment inputs and severe ice conditions during some years (Bustard and Schell 2002). These conditions can be greatly exacerbated with increased forest development in the area. As such development plans need to take into account the significance and fragile nature of these critical habitats.

90000000000

3.2 Chinook Salmon

3.2.1 Population Status and Trends

Within the Skeena watershed, the Morice is one of the most important rivers for chinook. The Morice River has produced on average 25% of the Skeena River escapements since 1950. From the early 1950's to 1985, chinook escapements were generally less than 5,000 spawners; however, since the mid-1980's, numbers have reached near record levels, with returns up to 15,000 fish. This rebound in escapement levels was reflected in the Skeena as a whole where escapements were low through the sixties to early eighties, followed by increased numbers in the last sixteen years (Bustard and Schell 2002).

Morice river chinook runs form an integral part of the in-river sport fishery and the Moricetown food fishery. Wet'suwet'en catch records originating from 1930 indicate an average annual chinook catch at Moricetown of 2,000 to 5,000 fish (Bustard and Schell 2002).

3.2.2 Limiting Factors to Production

Chinook production in the Morice system is probably limited by factors associated with spawning and rearing (Bustard and Schell 2002). Spawning and rearing limitations may include the effects associated with major floods, low winter flows, dewatering, extreme cold, redd superimposition and spawning gravel quality and quantity.

Low winter flows, extreme cold periods and freezing of redds in the upper Morice during some years can be significant factor in chinook survival. A combination of high fall flows that push spawners to the river edge and low late winter incubation flows is expected to lead to poorer chinook egg-to-fry survival in the upper Morice River (Bustard and Schell 2002).

Because spawning occurs over a period of roughly one month and chinook spawners tend to concentrate at sites offering the best spawning, the potential for chinook redd superimposition in the Morice is high (Bustard and Schell 2002). Reduced survival is clearly linked with chinook spawner abundance (Groot and Margolis 1991).

3.2.3 Habitat Condition and Protection

Prime chinook habitat within the watershed is found in the mainstem (spawning) and side channels (rearing) of the Morice River. Of critical importance is the four kilometer stretch of river downstream of the mouth of Morice Lake, with additional key areas downstream to the Lamprey Creek confluence. These habitats provide 97% of chinook spawning habitat in the Morice (Bustard and Schell 2002). Suitable gravel recruitment into these primary chinook spawning areas is probably limited, effectively determining how much spawning habitat is available. Generally, spawning site exposure to sediment events is expected to be rare because the Morice River is lake headed. However, forest management practices in the small watersheds flowing directly into key chinook spawning areas have the potential to introduce sediments and reduce the suitability of these sites (Bustard and Schell 2002).

3.3 Coho Salmon

3.3.1 Population Status and Trends

Since 1950, the Morice coho stock has accounted for an average of 6% of the Skeena watershed run. Morice coho returns have been volatile from the 1950's through to the 1990's with numbers dropping from as high as 15,000 fish in 1956 to 725 fish in 1991. Due to these extremely low escapements, the Department of Fisheries and Oceans declared a "coho crisis" during the late 1990's, initiating strong conservation measures to protect upper Skeena River coho stocks. Through restricting sport and Canadian commercial fisheries (responsible for approximately 65% of the harvest), escapements improved dramatically with estimated coho returns of 6500 fish in 1997 and 19,907 in 2000 (Bustard and Schell 2002). The sport fishery has recently been reopened, as populations appear to be increasing (Bustard and Schell 2002).

Wet'suwet'en fisheries also participated in coho conservation. Records indicate a drop in average catch from approximately 1500 fish per year prior to 1997 to less than 100 coho annually since then for the dip-net-fishery at Moricetown.

3.3.2 Limiting Factors of Production

Adequate spawner recruitment in the past has limited coho production. Poor ocean survival combined with high exploitation rates led to severely depressed populations for 30 years. The recent increased escapements indicate the high capability of the Morice watershed to support juvenile coho. Efforts to sustain coho populations in the Morice are dependent on ensuring strong escapements through fisheries management decision (Bustard and Schell 2002).

In late October and early November, floods due to rain on snow events help coho navigate beaver dams to reach spawning sites. However, fall freshets may also result in the scouring of redds (Bustard and Schell 2002).

Of the salmon species, coho rely on small tributaries for rearing the most. Weather conditions in both early and late summer determine coho distribution in non-spawning streams, with late summer flows and temperatures directly affecting the suitability of many coho rearing habitats (Bustard and Schell 2002).

Overwinter rearing mortalities in side channels and small tributaries are a result of poor water quality due to low flows, dewatering as flows decline during late winter, and predation as ice cover on the channels melts during the spring.

3.3.3 Habitat Condition and Protection

Coho are widespread throughout the Morice watershed, primarily using small streams. Therefore, habitat protection is an important issue for this species.

Habitat protection begins with accurate identification of coho habitat. During the decades of depressed coho abundance, many coho rearing streams may not have been identified by stream inventories conducted throughout the Morice watershed due to low stock densities. As such, known coho distribution may not be accurate.

The redistribution of newly emerged coho fry and yearlings into non-spawning streams and offchannel ponds is greatly influenced by road stream crossing structures. For instance, juvenile coho have limited ability to move upstream through road culverts (Bustard and Schell 2002).

The reliance of coho on smaller tributaries and off-channel ponds makes riparian management in and around these areas crucial. The maintenance of suitable water temperatures and flows as well as the reliable recruitment of debris for cover are important factors for coho rearing.

3.4 Pink Salmon

3.4.1 Population Status and Trends

Numbers of pink salmon returning to the Morice River have increased from approximately 5,000 in the early 1950's to 800,000 in 1991. Through the mid-eighties and early nineties, returns were consistently over 50,000 fish, but since 1993, annual pink spawner numbers in the Morice have fluctuated from 5,000 to 175,000. Morice pinks make up 9% of the Skeena River pink stock (Bustard and Schell 2002).

The Wet'suwet'en food fishery at Moricetown does not target pink salmon. However, the large escapements in the last 15 years have resulted in an average harvest of 3000-4000 pinks per year by the first nation. The sport fishery angling regulations have allowed the retention of two pink salmon per day in the Bulkley River and the lower two kilometers of the Morice River since the mid-1990's (Bustard and Schell 2002).

3.4.2 Limiting Factors on Production

Freshwater survival of pink salmon from egg to alevin, even in highly productive streams, commonly only reaches 10-20% (Groot and Margolis 1991). Primary factors affecting freshwater survival of pinks in the Morice are dissolved oxygen concentrations, stability of spawning beds, and freezing of redds (Bustard and Schell 2002). Redd sites with no surface flows tend to have low dissolved oxygen levels. In mid-winter, discharge levels decline in the Morice, resulting in reduced subsurface dissolved oxygen levels, as groundwater inputs comprise a larger portion of the discharge (Bustard and Schell 2002).

Climatic conditions such as winter severity and discharge levels can greatly influence pink salmon survival in the Morice River. Low discharge during the early winter can lead to direct freezing of redds in some key spawning habitats in Morice side channels in years with very cold temperatures and low snow cover. At the other extreme, rain on snow events can lead to extreme freshets after the pink spawning period, which can cause gravel shifting resulting in poor survival (Bustard and Schell 2002).

Predation of emerging pink fry by birds, coho smolts and resident char can be a significant factor in pink survival in the Morice watershed (Bustard and Schell 2002).

3.4.3 Habitat Condition and Protection

Increased forest development within the Thautil and Gosnell watersheds which empty into the main pink salmon spawning area of the Morice mainstem could lead to increased sediment

loading and reduced egg survival. As such, limited harvest rates and appropriate riparian management strategies in these areas are essential in order to protect critical downstream habitat.

3.5 Steelhead

3.5.1 Population Status and Trends

The Morice River is one of the most important steelhead systems in the Skeena drainage. The Morice watershed was estimated to have a capacity of just over 6,000 adults, or approximately 8% of the Skeena's potential of 80,000 steelhead. Mark-recapture surveys have estimated recent Morice steelhead escapements to be between 3300 (Lough 1995) and 6750 fish (Mitchell 2001).

A catch and release regulation has been imposed on the steelhead sport fishery in the Morice since the early 1990's. The Wet'suwet'en food fishery records at Moricetown indicate an annual catch of about 500 fish per year since the early 1980's (Bustard and Schell 2002).

3.5.2 Limiting Factors for Production

Adequate spawner recruitment into the Morice watershed has been greatly impacted by both the Alaskan and Canadian mixed stock fishery, the native fishery and hooking mortalities associated with the sport fishery catch and release regulation. These factors account for a 60% exploitation rate during some years (Ward et al. 1995). Recent observations also suggest that changes in ocean habitats have affected survival rates of steelhead (Bustard and Schell 2002).

Given adequate spawner recruitment, the limiting factor for steelhead production in the Morice is juvenile rearing habitat. Steelhead populations in the Morice are influenced by density control factors in some tributaries, as well as environmental extremes such as severe ice, low winter flows and freshets (Bustard and Schell 2002). The availability of steelhead rearing areas is restricted by low summer flows in Lamprey and Owen creeks, which may have a large influence on potential production of fry and parr. Water temperature in these systems is critical as well. Small increases in water temperature are expected to favour species other than steelhead, resulting in increased competition for similar habitats in these streams (Bustard and Schell, 2002). Rearing steelhead can be significantly impacted during severe winter conditions. As flows decline during the winter and the wetted area of the channels is reduced, standing water is created, water quality deteriorates, and predation by birds results in poor survival of steelhead parr. During the spring and fall, steelhead are subjected to high flows in the Morice watershed during the snowmelt freshet. These conditions are thought to result in the displacement of steelhead fry and small yearlings unable to find suitable refuge (Bustard and Schell 2002).

3.5.3 Habitat Condition and Protection

Some of the key steelhead habitats in the Morice watershed are subject to low summer flows, potentially high temperatures and high erosion potential. These conditions make steelhead habitats sensitive to forest development activities. Poor logging and road building practices in the Owen and Lamprey watersheds has resulted in significant sediment inputs into these systems including the headwater areas of core steelhead-producing tributaries such as Pimpernel Creek. Protecting the riparian areas and maintaining water temperatures in these systems is critical as well. A minor increase in water temperatures in both Lamprey and Owen could make these systems unsuitable for juvenile steelhead (Bustard and Schell 2002).

99999

3.6 Rainbow Trout

3.6.1 Population Status and Trends

The status of both the Morice Lake and Nanika-Kidprice lakes rainbow trout stocks are largely unknown. Creel survey data from October 1979 indicated rainbow trout were the most common sport fish captured in Morice Lake (Envirocon 1984), comprising 58% of the catch. Based on an additional tagging study, Envirocon (1984) estimated the rainbow trout population (individuals larger than 25cm fork length) in Morice Lake to be 4000-7000 fish. However, this estimate is expected to be high since it assumed no spawning moralities. No reliable estimates of the rainbow populations in Nanika-Kidprice lakes are available (Bustard and Schell 2002).

3.6.2 Limiting Factors of Production

Competition between steelhead and rainbow may be quite high since habitat requirements of juveniles are so similar. Resident rainbow are rarely noted at locations where significant steelhead populations occur (Bustard and Schell 2002).

Low nutrient levels in Morice Lake may be responsible for the slow-growing and late maturing nature of Morice Lake rainbow trout. As well the absence of lakes at the headwaters of Houston Tommy and Fenton Creek is suspected to be responsible for the small size (approximately 20cm) of the rainbow in those systems (Bustard and Schell 2002).

3.6.3 Habitat Condition and Protection

Rainbow trout populations in the Morice are primarily found in smaller streams as opposed to the mainstem Morice. The most important rainbow rearing habitats are found in the Nanika River. Environmental extremes such as dewatering and freezing of habitat, as well as displacement caused by flooding, may impact juvenile recruitment into Morice Lake and ultimately influence the rainbow population (Bustard and Schell 2002).

3.7 Cutthroat Trout

3.7.1 Population Status and Trends

Stock status of Morice River Cutthroat is unknown.

3.7.2 Habitat Condition and Protection

Cutthroat trout utilize small streams throughout the Morice watershed, and as such, are particularly susceptible to the impacts of forest development. Habitat issues for Morice River cutthroat trout stock are primarily linked to poorly installed road stream crossing structures that prevent upstream movements by juveniles and poor riparian management practices which lead to changes in hydrological flow and increased stream temperatures (Bustard and Schell 2002). Road development and access management of smaller lakes, streams and ponds should be an important factor in forest development planning in the Morice watershed.

3.8 Bull Trout

3.8.1 Population Status and Trends

British Columbia bull trout are designated as a blue-listed species, meaning populations are considered vulnerable or at risk. The current stock status of Morice River bull trout populations is largely unknown. Due to the difficulty associated with conducting bull trout counts, many studies have resorted to creel surveys or redd counts to provide estimates of bull trout populations. Such studies have estimated that the Morice River bull trout population may be limited to less than 1000 spawners (Bustard and Schell 2002).

3.8.2 Limiting Factors for Production

Bull trout prefer cold streams. Distribution of Morice River bull trout populations is strongly correlated with the coldest streams in the watershed. Temperatures rarely exceed 10 degrees Celsius in spawning sites and 12 degrees Celsius in juvenile rearing areas (Bahr 2002). Because bull trout are better adapted to cold water, they have a competitive advantage against other species in these areas. Conversely, bull trout are less effective competitors in warmer stream reaches.

The stability of redd sites is an issue in systems which are naturally dynamic (e.g. Crystal and Glacier creeks). Late fall freshets in bull trout spawning areas can lead to poor fry survival.

Fry displacement as a result of late freshets could be a concern. Side channel locations buffered from the full freshet flows are expected to be important refuge areas for bull trout fry (Bustard and Schell 2002).

Morice bull trout adults spend a significant amount of their time in areas easily accessed by anglers. Studies in other rivers with good angler access indicate that sport fishing can have a major influence on bull trout populations (Bustard and Schell 2002). In the Morice watershed, the Thautil-Gosnell confluence and the Nanika falls are thought to be particularly vulnerable to increased angler pressure (Bahr 2002; Bustard and Schell 2002).

3.8.3 Habitat Condition and Protection

Bull trout habitat protection issues are primarily linked to the impacts associated with forest development activities in the Morice watershed. Riparian management associated with small non-fish bearing streams flowing into bull trout spawning areas is important in terms of maintaining cold stream temperatures. The interception of groundwater and small seepage flows by roads can potentially affect bull trout spawning areas. In addition, channel stability in bull trout spawning streams can be greatly influenced by road stream crossings and the introduction of debris and course sediments. Finally, legal and illegal angler harvest of bull trout is greatly aided by increased access to key bull trout habitats (Bustard and Schell 2002).

1000000000

3.9 Dolly Varden

3.9.1 Population Status and Trends

The current stock status of Morice River Dolly Varden is unknown. There have been no direct studies on Dolly Varden populations in the Morice system. It is expected that Dolly Varden achieve relatively low densities in most stream systems, and that stocks are not threatened or declining in the Morice watershed (Bustard and Schell 2002).

3.9.2 Habitat Condition and Protection

Dolly Varden abundance, distribution and habitat preference throughout the Morice watershed, in conjunction with their small size, can lead to habitat protection issues that are linked to forest development activities. Dolly Varden are present in high gradient streams with marginal habitat. As such, careful attention to riparian management strategies aimed at protecting upstream sites from sediment, hydrological changes leading to low flow and peak flow impacts, and increased water temperatures is important. Although difficult, the proper installation of road stream crossing structures on these streams is important in order to allow fish passage. Drainage changes due to road construction and timber harvesting can also lead to impacts to key seepage habitats (Bustard and Schell 2002).

3.10 Lake Trout

Lake trout are known to be present in four lakes within the Morice watershed: Morice, Owen, McBride and Atna lakes (Envirocon 1984; Bustard and Schell 2002). The current stock status of Morice watershed lake trout is unknown.

3.11 Kokanee

The presence of kokanee in the Morice watershed has only been recorded in two lakes: Morice and Shea lakes. Based on a 1979 creel survey, Kokanee numbers are low in Morice Lake and only make up 1% of the total angler catch (Envirocon 1984).

3.12 Whitefish

According to the FISS database, three species of whitefish are present in the Morice watershed (Applied Ecosystem Management Ltd. 2001). Pygmy whitefish are reported in Owen and Morice lakes, and lake whitefish have been sampled in McBride and Morice lakes. Mountain whitefish is the most widely distributed and abundant whitefish species in the Morice watershed. Stock status of all three species is unknown (Bustard and Schell 2002).

3.13 Other species

3.13.1 Burbot

Burbot have been recorded in McBride, Morice and Owen lakes. However, no life history, stock status or distribution information is available (Applied Ecosystem Management Ltd. 2001).

3.13.2 Lake Chub

Within the Morice Timber Supply Area (TSA), this species has been identified as regionally important by the Ministry of Water Land and Air Protection (WLAP). Lake chub have been recorded in the Gosnell, Thautil, Lamprey and Owen creeks, and in Morice and Owen lakes (Applied Ecosystem Management Ltd. 2001).

3.13.3 Longnose Sucker

Longnose suckers inhabit McBride, Tagit, Owen and Lamprey creeks and their associated lakes. They are also present throughout the Nanika-Kidprice lakes system (Bustard and Schell 2002).

3.13.4 Largescale Suckers

This species is common throughout the mainstem of both the Bulkley and Morice rivers. In the early 1980's, largescale suckers made up 6% of snorkel observations of resident fish in the upper Morice River (Envirocon 1984). The Morice River between Gosnell and Owen creeks provides overwintering habitat for approximately 800 largescale suckers as counted during an aerial survey in late November 1979 (Bustard and Schell 2002). Largescale suckers have also been found in Collins, McBride, Morice and Owen lakes (Applied Ecosystem Management Ltd. 2001).

3.13.5 Redside Shiners

Redside shiners have been identified in Tagit creek and Owen, Collins, McBride and Morice lakes (Applied Ecosystem Management Ltd. 2001).

3.13.6 Longnose Dace

This species is abundant throughout the mainstem Morice River and Lamprey, Owen and McBride creeks. In 1979, longnose dace comprised 9% of the mainstem Morice River sample; however in subsequent years, this percentage decreased (Envirocon 1984). Dace are also found in the Nanika watershed.

3.13.7 Prickly Sculpin

This species is present in Morice and Nanika rivers and lower McBride and Lamprey creeks. They are also found in Owen and Gosnell creeks (Bustard and Schell 2002).

3.13.8 Pacific Lamprey

Pacific lamprey are abundant and widespread in the Morice watershed. Large numbers of pacific lamprey have been observed spawning in Owen and Lamprey creeks during June and July, and in the Morice River mainstem during late July (Bustard and Schell 2002).

The FISS database summarizes reports of peamouth chub and white suckers. Triton (2000) reports the presence of coast range sculpins and northern pike minnows in a reconnaissance level inventory (Bustard and Schell 2002).

9999999999

4.0 FACTORS AFFECTING MORICE FISH POPULATIONS

In the opinion of Bustard and Schell (2002), the primary factors influencing the health of Morice River fish populations are high commercial and sport fishery exploitation levels. In general, the integrity of fish habitat within the Morice watershed is sound; however, the cumulative effects associated with progressive forest road building and timber harvesting activities could have significant future impacts on Morice watershed fish habitat and its productive capacity.

4.1 Commercial and In-river Fishing

Based on existing information and population trends for Morice fish stocks, it is clear that commercial interceptions, and in some years, the in-river-fisheries, have had a severe impact on salmon and steelhead populations in the Morice (Bustard and Schell 2002). Management strategies aimed at curtailing the commercial fishery over the past decade have allowed increased spawner recruitment and provided evidence of the impressive habitat capabilities of the Morice watershed to produce native fish stocks. Recent escapement numbers for Morice coho chinook and steelhead provide a clear insight into these potential production levels (Bustard and Schell 2002).

4.2 Land Use Activities

Present land use in the Morice watershed is dominated by forest management activities, with minor amounts of grazing and mineral exploration and development. Forest management activities in most valleys have included progressive road development and timber harvesting utilizing the clearcut silviculture system.

According to the results of a baseline land use analysis for the Morice drainage, the most heavily developed sub-watersheds, as of the mid- to late 1990's, are located in the mid- and lower parts of the watershed (Figure 4). Owen, McBride, Lamprey watersheds and an unnamed watershed (Unnamed 26), showed the highest relative forestry impacts (Appendix B). The Morice mainstem (including first and second order tributaries) and Gold and Knapper watersheds scored moderately high in terms of relative land use.

Unfortunately the best available data for use in the analysis was dated 1995-1998. Therefore, the intention of this land use analysis is to provide a baseline data set in which to compare current land use data when it becomes available. This will allow for the identification of trends and possibly geographic areas, in which specific management objectives and strategies can be developed, implemented and monitored.

4.3 Fish Farms

Recent initiatives to establish fish farms on the north coast need to be closely examined in terms of the potential impacts to Morice native fish stocks. As outlined in Gottesfeld et al. (2002), the introduction of foreign species such as Atlantic salmon into the Skeena watershed could threaten the integrity of Morice steelhead and other native salmon stocks.




Figure 4: Relative impacts of land use within the Morice River watershed.

5.0 PRIORITY ISSUES

5.1 Morice Watershed Issues

Issues can be defined as problems, concerns or opportunities. Prior to determining objectives for any plan, the issues need to be identified. For the purpose of the Morice WFSP, issues were collated from a number of sources:

- 1. Summary reports produced for WFSP
 - Conserving Morice Watershed Fish Populations and Their Habitat (Bustard and Schell 2002);
 - Conserving Skeena Fish Populations and Their Habitat (Gottesfeld et al. 2002); and
 - Phase II Watershed-based Fish Sustainability Planning: Synopsis of key fisheries resource issues (Croft and Bahr 2002).
- 2. A previous watershed plan
 - Healthy Watersheds, Healthy Communities: Bulkley Morice Salmonid Preservation Group Draft Strategic Plan – Phase 1 (Tamblyn and Donas 2001)
- 3. The WFSP technical committee
- 4. The public via the Morice LRMP Fish Sector

Issues were organized into three categories:

- Fisheries management;
- Habitat (includes instream, riparian, upland and hydrological issues), and
- · Data gaps.

General or broad issues affecting many fish species or the aquatic ecosystem as a whole are listed in Appendix C1. Issues identified specifically for (a) species of fish are compiled in Appendix C2.

5.2 Selection of Priority Issues

Due to time limitations, the technical committee was unable to meet to prioritize issues. Instead, the authors have summarized the key issues as noted by Bustard and Schell (2002) in their conclusion and recommendation sections. There will be opportunities to further assess priority issues when the Morice WFSP resumes. A possible scoring matrix to help rank issues is in Appendix D.

Priority fish species as determined by Bustard and Schell are listed in Table 3.

Table 3. Priority fish species and stocks of in	erest
---	-------

Species / Stock	Reason for Priority Status
Nanika Sockeye	Unexplained population fluctuations dominated by low escapements since the 1950s.
Bull Trout	Blue listed species particularly sensitive to habitat degradation.
Steelhead	Vulnerable to fisheries management decisions and habitat impacts.
Coho	Vulnerable to fisheries management decisions and habitat impacts.

5.2.1 Priority Fisheries Management Issues

The issues in this and the following sections were identified as significant in Bustard and Schell (2002) but have not yet been ranked.

Significant fisheries management issues include:

- Lack of adequate coho spawner recruitment to the Morice watershed:
 - High Canadian commercial fishery bycatch in mixed stock fishery
 – sockeye, coho, steelhead, chinook; and
 - Heavy Alaskan commercial fishery harvest / bycatch steelhead, coho;
- Generous angling regulations for rainbow trout, lake trout and bull trout given lack of stock status information and oligotrophic nature of the system; and
- Past high First Nation in-river harvest rates for sockeye and steelhead during some years.

Addendum to:

A Plan to Conserve and Protect Morice Watershed Fish Populations and their Habitat – Stage II (Tamblyn and Croft, 2003)

Following the distribution of the final draft of *A Plan to Conserve and Protect Morice Watershed Fish Populations and their Habitat – Stage II* (Tamblyn and Croft, September 2003), Community Futures Development Corporation of Nadina received feedback we felt was important to pass to people receiving copies of the report.

Concerns were expressed over the wording of the following statement found in Section 5.2.1 Priority Fisheries Management Issues on page 24:

"Past high First Nation in-river harvest rates for sockeye and steelhead during some years"

Comments received:

- This statement should not be considered a priority or significant fishery issue.
- First Nations have a traditional right to harvest fish.
- The statement may lead to misconceptions by certain members of the public about the numbers of fish caught by the Wet'suwet'en.
- There is no current conservation concern regarding Bulkley steelhead populations.

We would like to point out that Wet'suwet'en Fisheries have been leaders in local stock assessment efforts over the past several years. They have run, or been partners in, projects to improve fisheries management in the Bulkley and Morice watersheds including coho, steelhead and sockeye mark-recapture programs. Currently, the Wet'suwet'en have voluntarily stopped harvesting sockeye at Moricetown in order to monitor stocks and allow the Maxan and Nanika sockeye populations to rebuild.

The Wet'suwet'en have also been a valuable member of the Bulkley Morice Salmonid Preservation Group and have participated in Strategic Planning for this group and are members of both the Morice WFSP planning and technical committees.

Sincerely,

Greg Tamblyn WFSP Planner

They Lambly

5.2.2 Priority Habitat Issues

Significant freshwater habitat issues include:

- Increased water temperatures, especially for bull trout and steelhead (Owen and Lamprey creeks);
- Road stream crossing barriers to fish migration, especially for Dolly Varden, cutthroat trout and coho;
- Loss of riparian integrity and function on small creeks (especially for Dolly Varden, cutthroat trout);
- Increased sedimentation and debris loading (particularly from the Thautil watershed) as a result of industrial development in all key habitats in the Morice watershed:
 - Upper Nanika River
 - Upper Morice River (Morice Lake to Gosnell Creek)
 - Mid-Reach Morice River (Gosnell to Owen Creek floodplain)
 - Key Steelhead Tributaries
 - Owen Creek
 - Lamprey Creek
 - Shea Creek up to falls
 - Upper Thautil River above Starr Creek
 - Key Coho Tributaries
 - Gosnell Creek floodplain and tributaries upstream from Shea Creek
 - McBride and Owen creeks
 - Lower accessible ends of smaller tributaries along the mainstem Morice and Nanika rivers
 - Key Bull Trout Tributaries
 - Upper reaches of Gosnell Creek
 - Denys, Loljuh and upper Starr creeks in the Thautil watershed
 - Glacier Creek
 - Houston Tommy and Gold creeks
 - Nanika-Kidprice Lakes
 - Des Creek and Lake.

6.0 OBJECTIVES AND STRATEGIES FOR SELECTED PRIORITY ISSUES

We have chosen two fisheries management issues and two habitat issues for which to write draft objectives and strategies. Draft targets are imbedded in some objectives as examples. Determination of final targets will require extensive efforts by the technical committee and are not possible at this time.

Please note that the WFSP technical committee and management agencies have not agreed to the following draft objectives, strategies and indicators. Their purpose is to promote dialogue once the Morice WFSP process resumes.

Issue 1: Lack of adequate coho spawner recruitment to the Morice watershed.

Lack of adequate coho spawner recruitment to the Morice watershed has been a major limitation to coho production in the system (Bustard and Schell 2002). Poor ocean survival combined with high exploitation rates by both the Canadian and Alaskan commercial fisheries have led to low escapements for much of the past 30 years. Since 1998, Fisheries and Oceans Canada has taken dramatic steps to protect Skeena River coho stocks by limiting recreational, First Nation and commercial coho fisheries.

The authors of this plan acknowledge that Fisheries and Oceans fish management biologists need to contribute to future iterations of these objectives and strategies.

Draft Objective

1.1 Increase average escapements of coho to the Morice watershed to <u>(a target #)</u> by <u>(year)</u> in order to fully reach productive capacity of the watershed.

Draft Indicator for Objective 1.1

Number of coho reaching the Morice watershed annually.

Possible Strategies / Management options

Fisheries Management

- Encourage terminal fisheries³ for sockeye in key locations in the Skeena watershed.
- Continue to adjust timing of commercial fisheries to allow maximum escapement of Morice coho.
- Encourage the use of selective fishing gear in the commercial and First Nations fisheries.
- Negotiate with the Alaskan commercial fishery to reduce catches of Bulkley bound coho.
- Restrict retention of coho salmon by recreational anglers should populations dip below (a number).

Research to fill data gaps

- Determine the carrying capacity and seeding requirements for the Morice watershed so that target escapements can be determined.
- Continue operation of the Toboggan Creek hatchery to provide an indicator stock for commercial fishery harvest levels.
- Develop improved in-season stock estimates for coho.
- Continue refining the mark-recapture program conducted at Moricetown Canyon.

³ A terminal fishery is the harvesting of adult fish at a location near, or at the end, of a stock's spawning migration.

Issue 2: Generous angling regulations for bull trout are putting bull trout stocks at risk given lack of stock status information and oligotrophic nature of the system.

The authors of this document acknowledge that Ministry of Water, Land and Air Protection biologists need to contribute to future iterations of these objectives and strategies.

Draft Objective

2.1 Reduce number of bull trout removed from the Morice River by anglers by an average of 50% from 2004 to 2013 or until population estimates and stock status are determined.

Draft Indicator for Objective 2.1

Average number of bull trout caught annually by anglers

Possible Strategies / Management options

Regulations

- Work with the Ministry of Water, Land and Air Protection to modify sport fishing regulations to reflect uncertainty in the bull trout populations in the Morice watershed.
 - Establish restrictions on angling at staging and holding areas for bull trout.
 - Reduce daily catch quotas and review size limits for bull trout.
 - Educate sport fishers about identifying bull trout vs. Dolly Varden by including information in regulations (synopsis).

Road access

 Restrict the construction of roads within 500 m of major staging and holding areas for bull trout.

Research to fill data gaps

- Monitor both adults and juveniles to assess bull trout populations over time by
 establishing long-term index sites for juveniles and redd counts on key spawning
 reaches in the upper Gosnell, Denys, upper Starr, and Glacier creeks.
- Conduct a multi-year creel survey at fishing locations along the Morice River to determine numbers of bull trout caught and if they are targeted or bycatch.
- Monitor impacts of angling on adult bull trout numbers in the Morice River.

Education

- Place signs at major fishing locations on the Morice River indicating what bull trout look like and identifying concerns about bull trout.
- Place the same information in the Freshwater Fishing Regulations for the Morice River.
- Promote release of fish caught in the Morice River that look like Dolly Varden.

Issue 3: High in-stream temperatures leading to lower fecundity, stress or mortality in salmonids, and improved ability to compete by minnow species.

Bull trout and steelhead are two species particularly susceptible to increased temperatures.

Draft Objectives

3.1 Maintain water temperatures in streams utilized by salmonids at temperatures below 12°C for bull trout and below 15°C for other salmonid species. (This objective could be modified to incorporate water temperature tolerance ranges for various life stages for each fish species of concern. Also important to consider is the temperatures at which non-salmonids start to out compete salmonids).

Draft Indicator for Objective 3.1

Peak temperatures (or mean daily temperatures) in streams utilized by bull trout and / or other species.

Possible Strategies / Management options

Industrial development

- During industrial development activities, ensure adequate riparian vegetation is left along S2-S6 streams and wetlands in watersheds sensitive to temperature increases including Owen and Lamprey creeks.
- Minimize modification of groundwater hydrology including springs and seepage areas when building roads and determining maximum allowable equivalent clearcut area to ensure adequate cool groundwater recharge to streams in low flow periods.
- Design and build roads to minimize temperature impacts on water (i.e. ditches collecting ground water and runoff are exposed to higher levels of solar radiation than buffered streams or groundwater).
- Minimize sediment run-off into waterways from roads and stream crossings, thus minimizing the water's ability to absorb solar radiation.
- Re-establish riparian zones where deemed appropriate alongside Owen and Lamprey creeks.

Research

- Continue with small stream temperature research to determine appropriate riparian zone widths for streams of various aspects as to avoid increasing instream temperatures.
- Monitor water temperatures annually in Owen and Lamprey creeks and other streams in which relatively small increases in temperature could change the relative composition of fish species.

Issue 4: Poorly installed road stream crossing structures impede juvenile and/or adult movements and migration.

Culverts can act as barriers to fish migration. Fish utilizing small streams including Dolly Varden, cutthroat trout and coho salmon are especially susceptible. Steep culverts act as velocity barriers, particularly for juveniles trying to access rearing habitat. Perched culverts can be barriers to adults migrating to spawning grounds and/or juveniles moving throughout a system.

Draft Objectives:

4.1 Ensure fish passage in fish bearing streams for juvenile and adult resident and anadromous fish species in 100% of new stream crossings.

Draft Indicator for Objective 4.1

Number of new stream crossing structures that inhibit upstream passage of fish based on best available determinants of fish passage.

Possible Strategies / Management options

- 4.1.1 Minimize number of stream crossings for roads.
- 4.1.2 Use options and technologies for road crossings that will ensure fish passage.
- 4.2 By 2008, restore uninhibited fish access to 20% of historically available habitat that is inaccessible in 2003 due to poor stream crossing structures.

Draft Indicator for Objective 4.2

Area $(m^2 \text{ or } km^2)$ of fish habitat to which full access has been restored compared with the area of historically available habitat.

Possible Strategies / Management options

Road Development

- 4.2.1 Replace existing structures that prevent upstream fish movement or those that impede access to certain habitats associated with specific life stages of fish present in the system.
- 4.2.2 Where access is not a priority for user groups, remove existing structures, permanently deactivate roads, and prevent motorized access.

Research

- 4.2.3 Complete a full inventory of road crossings in fish bearing streams to determine which crossing structures currently restrict fish passage or could possibly impede fish movement in the future.
- 4.2.4 Calculate net habitat "loss" by determining the quantity and quality of potential spawning, rearing and overwintering habitat, for all fish species, upstream of barriers caused by road stream crossing structures.

7.0 MONITORING FRAMEWORK

Because the Morice WFSP is still underway, developing a monitoring plan is not feasible at this point. The final monitoring strategy will be directed by the final objectives and indicators of the watershed plan. Below are some monitoring concepts and the key components of a monitoring plan.

7.1 What is Monitoring?

Monitoring is regular or ongoing systematic testing, sampling or tracking of specific parameters in order to collect information or to determine trends. A number of monitoring approached can be implemented through a strategic plan including baseline, implementation, effectiveness, project and compliance monitoring.

7.2 Why is Monitoring Important?

Monitoring is a vital component of any plan or project. It is required to determine the effectiveness of strategies and actions in meeting the objectives of the plan. In the case of the Morice WFSP, monitoring will be used to determine the progress made toward mitigating impacts on fish populations, fish habitat and the aquatic ecosystem as a whole.

Monitoring is also integral to improving a plan through adaptive management. Adaptive management uses well-designed monitoring programs to inform management decisions and allow adjustments as circumstances change or knowledge is gained.

7.3 Monitoring for the Morice WFSP

We recommend two main types of monitoring for the Morice WFSP:

- Implementation monitoring: Determine the status of the actions laid out in the implementation plan. We envision this monitoring to be done by the Implementation Committee who will actually put the plan into action. Their role would be to monitor the status of plan implementation and to address challenges and barriers to implementation.
- 2) Effectiveness monitoring: Determines to what extent objectives in the plan are being met. Effectiveness monitoring will help determine if the implemented management options or strategies have been successful, or perhaps whether or not objectives are feasible. Effectiveness monitoring is based on the indicators developed in the plan.

Effectiveness monitoring has three main stages according to Mackay (1998):

a) Baseline evaluation: Collating existing information for areas of concern, identifying data gaps and determining what potential impacts may arise from different land use and fisheries management decisions. This has been done to a large extent by Bustard and Schell (2002) and work completed by the Watershed Restoration Program (Ecofor 2001).

Morice Watershed-based Fish Sustainability Plan - Stage II


- b) Site Design: Selection of control and monitoring sites considering indicators and the types of information that will be required to determine if objectives are being met. Careful site selection to consider control or reference sites is vital to adaptive management.
- c) Measurements:
 - Precedence measurements: Measurements to fill data gaps or measurements prior to land use development or prior to the implementation of a strategy outlined in the plan.
 - Condition measurements: Measurements taken following land use or strategy implementation (i.e. results monitoring).

Much of the monitoring detailed in the final Morice WFSP is expected to be long-term. Long-term monitoring is required to track trends and to determine cumulative effects of land use on aquatic ecosystem integrity. In addition, some aspects of streams fluctuate naturally throughout a year or from year to year. To control for this natural variation, we recommend 1) monitoring over a range of geographic areas and 2) monitoring benchmark sites in an attempt to separate naturally occurring events from human induced impacts (e.g. temperature increases due to climate change as opposed to clearing of riparian vegetation).

A well-designed effectiveness monitoring plan should include (Mackay 1998):

- The monitoring objectives;
- Statistical considerations;
- What parameters to measure and the limits of acceptable change for each parameter;
- The benefits and advantages of monitoring selected parameters;
- Where monitoring will occur and why;
- How this plan can complement current monitoring activities;
- How frequently measurements are made;
- How the introduction of error and bias can be minimized (quality assurance and control);
- How the program's cost effectiveness can be maximized;
- How results are to be analyzed; interpreted and reported; and
- How long monitoring should continue in the absence of any exceeded limit of acceptable change.

Figure 5 shows a set of steps that could be taken to develop a monitoring plan.



Defined by the trade-off between cost, requirements to meet objectives and the need for statistical confidence

Figure 5: Possible steps to develop a monitoring plan. Source Mackay (1998) with minor modifications.

7.4 The Monitoring Toolbox

In monitoring, as in a mechanic's shop, there is no magic tool. A number of tools or indicators is required to evaluate the integrity of an aquatic ecosystem and the impacts of various resource management developments and decisions. The challenge is to find the most cost-effective set of tools to fill data gaps while gaining maximum feedback.

Mackay (1998) provides a number of options for monitoring the potential impacts of forest development on aquatic ecosystem integrity (Table 4).

Table 4. Possible tools for monitoring forest development (adapted from Mackay 1998)

Impact issue	Tools	Reconnaissance assessment tool?	Detailed assessment tool?
	turbidity	X	X
Sedimentation	Total suspended sediments	X	X
	Substrate cores		X
	Sediment traps		X
	Conductivity	X	
	Cobble embeddedness	X	
Channel Morphology	Pebble count	X	X
Chaimer Morphology	Bank stability	X	
	Large woody debris	X	
	% pools	X	X
	Degree of channel aggradation / degradation	X	
	Bankfull width and depth	X	
Water Quality	Temperature	X	X
Water Quains)	Fecal pathogens	X	
representation of the second	Sulphate	Translation with the pro-	X
	Dissolved oxygen	X	X
	Pesticides / Herbicides		X
Water Quantity and Timing	Bankfull width and depth	Х	
Tillinig	Peak flows		X
	Low flows		X
	Timing to peak flows		X
	General discharge curves	X	<u> </u>
	Groundwater flows		X
Stream productivity	Nutrients 1997	X	
	Alkalinity	X	
The state of the s	Chlorophyll a	X	
	Algae biomass	X	
	Periphyton	X	V
	Benthic invertebrates	X	X
	Fish	X	A

Monitoring the effects of fisheries management decisions is another component of the plan. Historically, stock assessment (numbers of fish, combined, in some cases, with weight, size and age measurements) has been the dominant tool:

- Adult counting fences;
- Redd counts;
- Juvenile index sites trapping / electroshocking;
- Catch per unit effort;
- Aerial, in-stream, stream shore counts for spawning adults;
- Mark / recapture;
- · Inter and intraspecific competition studies; and
- Creel surveys.

However, additional measurements providing greater insight into the health of fish could also be tracked:

- · Fecundity;
- Genetic diversity;
- Disease proneness; and
- Toxin levels.

Finally, an assessment of the social and economic value associated with fisheries management decisions could be used as an index (i.e. does an increase in escapement equal greater economic stability?).

7.5 Data Management

An effective monitoring program requires that collected data and results be accessible to the organizations responsible for implementing and reviewing the plan. On option is to utilize the Upper Skeena Atlas website as a depository for summarized information – reports, implementation committee meeting summaries, etc. The GIS portion of the website could be used to display monitoring locations, collected data and information about each sampling site. This website is part of the Community Mapping Network and the Sensitive Habitat Inventory Mapping project (http://www.shim.bc.ca/), run with the help of Brad Mason of Fisheries and Oceans Canada. The site integrates data from many sources including major government fisheries and watershed databases and makes it accessible through a user-friendly mapping system that everyone can access with minimal training.

8.0 IMPLEMENTATION FRAMEWORK

An implementation program specifies who will do what and when to put the plan into action. In other words:

- What agencies, organizations or companies are responsible for carrying out the various "actions" of the plan and when;
- Who is responsible for funding the various "actions" outlined in the plan and
- How the Morice WFSP will be integrated with related planning processes.

The implementation plan will also outline:

- The sequence and timeline for implementing activities;
- The creation of a data management system to monitor progress and maintain records; and
- A contingency plan to identify options should an organization be unable or unwilling to meet its designated responsibilities.

Potential groups involved in implementing aspects of the Morice WFSP include:

- Forest licencees: HFP, Canfor, BC Timber Sales branch of the Ministry of Forests;
- Ministry of Water, Land and Air Protection: Ecosystems Section, Fish and Wildlife Science and Allocation Section, Environmental Quality Section;
- Ministry of Sustainable Resource Management: Planning Section; Information Section;
- Fisheries and Oceans Canada various branches;
- First Nations;
- Tourism operators;
- · Community groups; and
- New industrial developers to the area.

Plans through which parts of the Morice WFSP could be implemented:

- Morice LRMP;
- Morice-Lakes Innovative Forest Practices Agreement;
- Sustainable Forest Management Plans; and
- Morice Sustainable Resource Management Plan

9.0 NEXT STEPS

The WFSP process, as outlined in MELP & DFO (2001) envisioned four stages. Stage II involves developing strategic direction (i.e. goals), management objectives and strategies. Stage III establishes an implementation and monitoring plan. The final stage, Stage IV, puts the plan into action, allowing opportunities to modify the plan through feedback derived from monitoring.

This document summarizes progress in Stage II of the process. Completing Stage II of the Morice WFSP will require:

- 1. Obtaining sufficient funding;
- 2. Confirming government, public and forest industry interest and support, and identifying available expertise to participate in and implement the Morice WFSP;

Morice Watershed-based Fish Sustainability Plan - Stage II

- 3. Completing a Local Strategic Overview that considers social, cultural, economic, political and ecological issues. The Morice WFSP process is in a unique position because the concurrent Morice Land and Resource Management Planning process has summarized most of this information;
- 4. Reviewing and ranking issues;
- 5. Developing objectives, indicators, targets, and strategies for key management issues; and
- 6. Assessing the advantages and disadvantages of each strategy / management option.

Steps in Stage III include:

- 1. Developing a monitoring and assessment program to evaluate performance of the plan and
- 2. Developing an implementation strategy required to achieve objectives, including projects, activities, manpower, and funding.

Given the current provincial government land use planning system, evolving government legislation and policy, and reduced government staffing levels, implementing the Morice WFSP is reliant on its integration into other planning processes. For a short time, opportunities exist to have portions of the Morice WFSP implemented through the Morice Land and Resource Management Plan. However, the Morice LRMP is scheduled to be finished by March 2004. If the Morice WFSP is delayed much longer, a tremendous opportunity to have WFSP recognized in a higher level plan will be missed.

10.0 REFERENCES CITED

- Applied Ecosystem Management Ltd. 2001. Morice Forest District fisheries data and information compilation project. Prepared for Ministry of Environment, Lands and Parks. Skeena region. Smithers, B.C..
- Bahr, M. 2002. Examination of bull trout (Salvelinus confluentus) in the Morice Watershed. Report prepared for Forest Renewal BC. University of Northern British Columbia.
- British Columbia Ministry of Environment, Lands and Parks and Fisheries and Oceans Canada. 2001. Watershed-based Fish Sustainability Planning. Conserving BC Fish Populations and their Habitat. A Guidebook for Participants.
- Bustard, D. 1999. 1998 stream inventory report Gosnell Creek Watershed. Report prepared for Forest Renewal B.C. (Northwood Inc.).
- Bustard, D. 1986. Some difference between coastal and interior stream ecosystems and the implications to juvenile fish production. *In*: Proceedings of the workshop on Habitat Improvements, Whistler, B.C. 8-10 May 1984. *Can. Tech. Report of Fisheries and Aquatic Sciences* No. 1483.
- Bustard, D. and C. Schell. 2002. Conservation of Morice Watershed Fish Populations and their Habitat. Stage II Biophysical Report. Morice Stage II Watershed-based Fish Sustainability Plan. Prepared for Community Futures Development Corporation of Nadina. Smithers, B.C..
- Croft C. and M. Bahr. 2002. Phase II Watershed-based Fish Sustainability Planning: Morice Watershed. A Synopsis of Fisheries Resource Issues. Prepared for Community Futures Development Corporation of Nadina. Smithers, B.C..
- Ecofor Consulting Ltd.. 2001. 2001 FRBC Watershed Restoration Program Interim Restoration Plans for the Buck, Houston Tommy, Kidprice, Morice Lake, Nadina & Owen Watershed Units within the Morice Forest District. Prepared for Houston Forest Products / Canfor-Houston Division.
- Envirocon Ltd. 1984. Physical and hydrological baseline information. Vol. 2. In:

 Environmental studies associated with the proposed Kemano Completion Hydroelectric development. Aluminum Company of Canada, Vancouver, B.C..
- Envirocon Ltd. 1984. Fisheries resource information of the Nanika River system: baseline information Vol. 3. *In: Environmental studies associated with the proposed Kemano Completion Hydroelectric development*. Aluminum Company of Canada, Vancouver, B.C..
- Envirocon Ltd. 1984. Fisheries resource information of the Morice River system: baseline information Vol. 4. In: Environmental studies associated with the proposed Kemano

- Completion Hydroelectric development. Aluminum Company of Canada, Vancouver, B.C..
- Gottesfeld, A.S., K. Rabnett, and P.E. Hall. 2002. Conserving Skeena Fish Populations and Their Habitat. Skeena Stage I Watershed-based Fish Sustainability Plan. Skeena Fisheries Commission. Hazelton, B.C..
- Groot, C. and L. Margolis. 1991. Pacific Salmon Life Histories. UBC Press, Vancouver, B.C.
- Horn, H. and G. Tamblyn. 2000. Morice Planning Area Background Report: An Overview of An Overview of Natural, Cultural, and Socio-Economic Features, Land Uses, and Resources Management. Produced for the Prince Rupert Interagency Management Committee. Smithers, B.C..
- Land and Water B.C. Inc. 2003. Data provided by Dwayne Meredith, Water Resource Specialist. Victoria, B.C..
- Lough, J. 1995 (Draft Report). Estimating the population of adult steelhead in the Morice Rive using mark-recapture methods, 1993/94. Skeena Fisheries Section. Smithers, B.C..
- Mackay, S. 1998. EWET: The Ecological Watershed Evaluation Toolbox Guidebook. Version 1.0. Prepared for: Pollution Prevention Section, BC Environment. Smithers, BC.
- Mitchell, S. 2001. A Peterson capture-recapture estimate of the steelhead population of the Bulkley/Morice river systems upstream of Moricetown Canyon during autumn, 2000, including synthesis with 1998 and 1999 results. Submitted to Steelhead Society of British Columbia, Bulkley Valley Branch. Smithers, B.C..
- Remington, D. 1996. Review and assessment of water quality in the Skeena River Watershed, British Columbia, 1995. Can. Data Rep. Fish. Aquat. Sci. 1003.
- Statistics Canada. 2003. 2001 data on Statistics Canada website: http://www.statcan.ca/start.html
- Tamblyn and Donas. 2001. Healthy Watersheds, Healthy Communities: Bulkley-Morice Salmonid Preservation Group Draft Strategic Plan Phase I. Community Futures Development Corporation of Nadina. Smithers, B.C..
- Triton Environmental Consultants Ltd. 2000. Reconnaissance (1:20,000) fish and fish habitat inventory in the Morice River Watershed. Prepared for Northwood Pulp and Timber Ltd. Houston, B.C.
- Ward, B.R., A.F. Tautz, S. Cox-Rogers, and R.S. Hooton. 1995. Migration timing and harvest rates of the steelhead trout populations of the Skeena River system. PSARC Working Paper S92-07.

APPENDIX A MEMBERS OF THE MORICE WFSP PLANNING AND TECHNICAL COMMITTEES.

Project Coordinator: Mary Swendson - CFDC Nadina / Greg Tamblyn (final month)

Planning Team

- Dana Atagi, WLAP (initial participation)
- Brenda Donas, DFO
- Martin Forbes / Dale Gueret, Fisheries and Oceans Canada (DFO)
- Allen Gottesfeld, Skeena Fisheries Commission
- Walter Joseph, Wet'suwet'en Fisheries
- Sharon Roberson, BC Federation of Fly Fishers / CFDC Nadina
- Greg Tamblyn, CFDC Nadina
- Gord Wadley, Bulkley Morice Salmonid Preservation Group (not active member)

Technical Team

- Gary Baptiste, Wet'suwet'en Fisheries
- Matt Jessop, Ministry of Sustainable Resource Management (early 2002)/ Ministry of Water, Land and Air Protection (2002 - March 2003)
- Vesna Konic, DFO (2002)
- Allen Gottesfeld, Skeena Fisheries Commission
- Tom Pendray, DFO
- Stefan Schug, Wet'suwet'en Fisheries
- Jim Schwab, Ministry of Forests (not active member)
- Greg Tamblyn, CFDC Nadina
- Melissa Todd, Houston Forest Products / IPFA
- Laurence Turney, IFPA (part of 2002)
- Carl Vandermark, Canfor / IFPA (part of 2002 / 2003)

APPENDIX B Morice WFSP Preliminary Land Use Analysis

460-600600-93600	のないとはいいのである	460-600600-32800	が 100 mm 大	460-600600-75800	#00-000000-00000	100 c00c00 35c00	460-600600-44500		460-600600-86900	を から	460-600600-86100-31000	460-600600-86100	TO COCCO OCAO	460-600-00-07 100	100 00000 07400	400-000000-01400-04100	460-600600-64400-70	460-600600-64400-59100	460-600600-64400-48600	460-600600-64400-43500	460-600600-64400-37000	460-600600-64400-11700	460-600600-64400	一次の 一般の 一般の 一般の	460-600600-67600-46	460-600600-67600	A. T. C.	460-600600-73500-82700	460-600600-73500-79700	460-600600-73500-69400	460-600600-73500-50000	460-600600-73500-27600	460-600600-73500-22600	460-600600-73500-20700	460-600600-73500	400-00000-00-00-00-00	460-600600-36400		460-600600-17000-80100	460-600600-17000-67900	460-600600-17000-6	460-600600-17000-50600	460-600600-17000-35200	460-600600-17000-27400	460-600600-17000		460-600600-50800-0		460-500600-50800-72200	460-600600-50800-51300	460-600600-50800-48000	460-600600-50800-3	460-600600-50800-17400	460-600600-50800	100-0000	400	100
Unnamed 27		Unnamed 26	を の	Unnamed 25	O mornio	I Innamed 24	Tagit	September 1	Unnamed 23		OU Unnamed 22	L	State Control of the	reduces	11642420	Cinding	1	OT Demend 20	L	L	L		L	の 一日 一日 一日 一日 日 日 日 日 日 日 日 日 日 日 日 日 日	L	L	2	2700 Unnamed 15	L	L				\perp	Atna		Dimpernell	The second	0100 Unnamed 8		L				Houston Tommy		-00500 Gosnell Creet	1	2200 Starr			L	7400 Unnamed 1	Thautil River		Morice River	NAME
1613	語なる時代	3351		390		PESE	42	· · · · · · · · · · · · · · · · · · ·	3658	100	-	1	40	3 0	3030	100 miles	2000	1	1	1	L	L	6	2		-	1	L	L	L	Ц		4	15	12	23	F 0			L	L	L	34		-	W.		42	92	1 4	1 1 1	41	16		100		Area
0.00	133	51 26.13	9	0.00	E I	3	0.00		0.00	1		T	3030 3.11	U	16	75.00	†	†	t	Τ.	9360 0.00	T	Ť	92	-	+	90	Т	2612 0.00	\vdash	Н	\forall	7	7	1		6781 16	5	0	✝	╁	۲	3414 0.	047 0.	,,	4		201	1	1968 6	+	T	-	12172 7.	120	힉	+
	M	0				- 100 mg		3	-	2	+	+	- 8		1		+	+	+	╁	╁	-	╀	37	7	╀	53	H	-	\vdash	Н	\dashv	\dashv	+	- 1	36	16.45	1	W.	╀	╀	┝	0.00 0	Н	\neg	8	П	<u>ند</u>	+	6 8 6	╁	╀	L	05 0	-	-	Road Road
0.00	8	Г	0		SA .	0.00		300	Т	0.00	8	3 8	0.10	0.3	17	3	3 8	8 8	200	0.15	0.00	0.38	9	0.00	0.00	8	0.00		8	0.00	0.00	0.00	0.00	0.0	7		0.08	0.04	8	0.00	8	0.00	8	.00	П	-		2	0.00	8 8	9.8	0.00	0.00	1 1	8	-	٢_
	7.00		189 80	H		1000		1 20 30	-	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	-	+	何一 少田 八世報		100 miles		+	+	+	+	-	\vdash	H	1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -		+	新国共产生	L	H	_		\parallel	+	+		5	+	100		\vdash	-	-	H	_	-	の記録	-		+	+	+	L	H	\parallel	3		Score loo
	の情報の表記を	6.783	発記に発力		1014	1010	2.503	影響の発表は		经营业 医耳动			語のない方を必然の	6.116	2 112	2000	0.00	0.000	0.00	0.000	0.000	11.508	8.703	O STATE OF		0.000	0	1	0.000	0.000	0.000	0.000	0.000	0.000	0.000	73	13.543	E0 E70	0.000	0.000	0.000	0.000	0.000	0.000	3.362	を の の の の の の の の の の の の の の の の の の の		9	0.043	0.00	6.571	0.000	0.000	2.255	00.000	90 003	Str length
0.0	16.1	16.1	0.0	0.0	22.5	18	3.5	257	0.0	175	0.0	0.00	00	0.7	15	0.0	0.0	0.0	0.0	0.0	0.0	15.8	1.1	0.00	0.0	0.0	0.00	0.0	0.0	0.0	0.0	0.0	0.0	0.0		21.7	43.7	OF A	6	0.0	0.0	0.0	0.0	0.0		0.4	0.4	1	0.0	200	13.4	0.0	0.0	0.8	0.0	Sec.	
2	1 7		0 1		1	3.0	Ÿ.	0.0		0.0	Т	\dagger	2	3	- 2	10	t	t	\dagger	\dagger	t	T	t	7	.0	t	00		r	r			1	1	_	7 9	\dagger	L . Gro	-	t					\neg			2	†	\dagger	t				8		+
200000000000000000000000000000000000000		4.543	100			100 May 100 Ma	n de la companya de l			S. See See See See See See See See See S			1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	0.200	0 255	35 13	0.00	0.00	1.020	20.31/	0.000	4.2/4	9.509	3000		0.00	0.0	0.0	0.000	0.000	0.000	0.000	0.000	0.000	1	45.5	5 350	10 134	0.000	0.000	0.000	0.000	0.000	0.000			2.914	5	0,000	3,007	0.000	0.000	0.000	2.727		03.037	Index Rd Length
0.00	0.14	0.14	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	3 8	000	0.0.	001	0.00	0.00	200	9 6	0.11	0.00	0.72	0.03	1		T			0.00	0.00	0.00	0.00	0.00		0.00		00.00	0.75	0.00	0.00	0.00	0.00	0.00	0.00			0.01		0.00	0.00	0.00	0.00	0.00	0.02	0.11	011	Rd Length
	3	Г	0 1	П	0	0		0 7	ŝ	0.000	1	\dagger	# (P#200 000			3	†	\dagger	†	\dagger	t	+	t	0	T	t	U en g	Г	r	H			1	1		7 8	\dagger	9		t	H	\vdash				H		1 8 1	†	†	\dagger	-	-	П	2	1	Score
		45	1000000		2000	6	9.	100		1000年の一日本の日本の			and the state of t		14	19						-	9	STATE OF THE PARTY.			The state of the s		0.0	0.0	0.0	0.0	0.0	0.0	0.0		28 55	030	0	0	0	0	0	0		· ·	31	Ž.	0	5 0	42	0	0	П			Area Logged w/in 30m (ha)
88		1.3	320	0.0	11	0.2	0.4	133	3	17.2		0.0	000	7.0	1000	7.0	0.0	0.0	0.0	0.0	0.0	1.9	0.2		0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0			12	24 0.7		0.0	0.0	0.0	0.0	0.0	0.2	armin and	0.1	***	0.0	0.0	2.3	0.0	0.0	0.1	0.0	0.80	% Log w/in
0.0	100	T	0.0	П	0.2	0.40		L. C.		0.0	-	t	4		A SCHOOL ST	4	\dagger	\dagger	\dagger	t	\dagger	\dagger	t	1.0		t	The state of the s		H	-		1	1	+		10	\dagger	0.3		H	-	-		+		0.1	T	2	\dagger	\dagger	+	H			0.8		Index
	はははは	1080				95	238	1		20 MAGE!		\dagger	And Charles and	17	212			,				14	47	遊戲	+	9	J.	-	0.0	0.0	0.0	0.0	9.0	0.0	+	100	416	50	0		0	0	0	0			1		3		37	0	0	\vdash	200 to	-	Area
	対応 発展 機	F				N. Carlotte		No. of Persons	-	100 Contraction of the Contracti		+	Sales of Section		1	1994	+	+	+	+	+	-	-	では、 できる			Street Live				0.0					116	7 -	888							6		578	918	+	+	0			8			
0.0	32.2	32.2	0.0	0.0	27	27	2	0.0	0.0	0.0	1	0 0	00	0.0	5.0		0.0	0.0	0.0	0.0	0.0	20.1	1.4	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	00	333	20.5	2.8		0.0	0.0	0.0	0.0	0.0	4.5		- 1	22	0.3	0.0	20.4	0.0	0.0	4.3	10.2	10.3	% watershed
	10		おき 公園	П				公司 (本)	Γ					9. 51	25										Т		京大学 大学						1	1	Special graphics	70	1	3	Г				1				200	2	1	1		П		\neg	2	╁	Index
0	11	11	可能够够	0		0	0	SAN SAN SAN	Г	0	0	0	Section 1	1	ONE		0	0	w	ē.	0	10	23	では、	0	0	Same of the same of	0	0	0	0	0	0	5	SERVE		8	A		0	0	0			3		3		0	0	0	0	0		1/4	1	# stream

В	9	SAS.		1000	28	公衛在 公司在		With John		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		のなると	-	100 Carlotte	C SALES OF STREET	1	1000 Harris			1	1	-	١		~17			1	1		1	8	ij.	١	١				١	1	١	1	2	1	0.00		1	١	1	١	T	1	13		10	8	1	077	-	1		١	- 1	1)37 0	00m Density
	6	0 16	0.00	3	0.14	0.00	0.00	0.00	0.00	0.00	0.00	שעים	0.00	0,00	2	3			0.01	0.5	0.00	0.00	0.00	0.01	0.11	0.00	0.12	0.03	0.00	200	3 5	3	0.0	8	000	0.00	0.00	0.00	0.00	0.00	3 5				0.26	0 15	38	3 8	3 8	3 8	3 8	3 5	01	0.01	100	100	3 8	0.03	3 8	8	8	8	.02		Ц	
	4	Carrier and Control				September 1		State of the last		(A) (A) (A) (A) (A)		(\$40)56 4 (2015)	•	Company Property				1		10				L					Section of the second			Section 1	Y.					L		1	1	Section of		1	400	2	1	1	1	1	1	1		数はない	Contract of the		1	1	4	1			Ц	2		Score w
139	作品があるというなど	205	A STATE OF			45	Printer and the second	Section of the sectio	o	E STATE OF THE STA	10	16		Control of the second second	e por de la companya			化水料水料	14	124	0	0	0	0	0	0	17	2	Particular Services				化 体系 在 1000 元	0.0	0.0	0.0	0.0	0.0	0.0	0 0	000	- 3	436	86	350	24	0	0	0	0	٥	0 !	24	のとのでは、		57	0	0	0	42	0	0	15		642	ਸ਼ੂ a)
1.4	0.1	1.0	0.0	0.0	13	1.3	0.0	13	0.2	0.2	0.4	0.4	0.0	- 1	0.0	0.0	0.0	1.0		0.1	0.0	0.0	0.0	0.0	0.0	0.0	9.1		- 18	0.0	00	- 1	0.0	0.0	0.0	0.0	0.0	0.0	000	0.00	000	- 13	20 x 20 20 20 20 20 20 20 20 20 20 20 20 20	1.3	- 12	200	0.0	0.0	0.0	0.0	0.0	0.0	- 1	0.1	- 1	7.0	0.0	0.0	0.0	2.3	0.0	0.0	0.1	0.8	0.8	30m of stream
	0 6	П	0 1831		3 7	_	0	_	2		() () ()	_		_	0 1 1 1 1 1			4 3		1 100 1-00			-	T		T	t	†		0	1		1 and 1 and		-		T	t	1	†	1	_	10	1	-		1	1	1	1	1	1		1	-		1	1	1	1	1			8 5	1	Score
3428	10000000000000000000000000000000000000	3540	· · · · · · · · · · · · · · · · · · ·		地は様々され	1080		Contractor Contractor		95	1. 100	238	を変する					高い という	213	1221	0	0	0	c	, ,	,	ì	1	477	Section 1	0.0	0.0	を放けた	0.0	0.0	0.0	0.0	9 6		3	00	0.0	5116	975	4141	686	0	0	0	0	0	0	686	ではいる	578	918	30	0	0	370	0	0	518	10000000000000000000000000000000000000	JACE	Logged
33.7	17.0	17.0	0.0	0.0	32.2	32.2	0.0	- 44	2.7	2.7	5.8	5.8	0.0	- 1	0.0	0.0		5.8	5.8	N.	0.0	0.0	0.0	0.0	0.0	0.0	20.1	30.1		0.0	0.0	0.0	0.0	0.0	0.0	0.0	200	0.00				0.0	ĝ.		26.5	1000	0.0	0.0	0.0	0.0	0.0	0.0		- Sept. 1.1	1.1	1	П	0.0	0.0	20.4	0.0	0.0	4.3	ZOL SECTION		Logged
	. 6 Sec		海祖		70		でを				2		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		1000			2	Γ	徳山 安治					T	Ī	T						新文·										7			李 3								7. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.										- The -	The second second	SCORE
8	67	67	0	0	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	===	一种工作的	0	を行うない	0	大量以吸作的	0	10.45	0	()	0	0	歌樂就是		100	0	0			3 9	2	0	3 10	23	は、意味な	0	0	是是	0	0		,			-	0	0	92	12	80	なると	0	0	0	0	0	0	з		2	93	0	ڻ.	0	0	0	0	000	174	1/4	X-ings
113.95	1000	261.46	533	19.86		42.04		13.52	- 大学を変え	55.47		71.99		79.88	0 149.7	40.41	109.34		61.87			54.08	80.32	22.667	230.00	20.02	250 62	75 71	808.90	60.6	24.10	36.56	0 548.15	16.12	31.38	35.70	24.00	20.51	85 24	57 05	45.86			102.57	234.93	3 525.1	25.00	11.08	23.97	46.99	80.75	20.96	316.44	2	1263.47	3 -1087.1	274.90	316.03	24.40	48.89	119.59	35.14	268.16		1486.54	Length
- A - A - A - A - A - A - A - A - A - A	0.26		0.00		0.26		0.00		0.00		0.00		0.00		5 0.00			0.02		0.05							1	1		0.00			5 0.00				1	1					0.27			0.01								0.00		10.0								0.12	9	of Stream
	6 7	1			. 6		1000		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	-	· 大學公		10 m		1					2.34			T	T		T	1		_								Ī	T	1	1			8											の変数を		2000								1		Score
27	- 32		6		40		- B		THE REAL PROPERTY.		10		6		6			1 10 2	-	-16		T	T	T	1	T	1	1	-6	8		_	100			T	T	T	1	1			45		-	8			1			-	-	**************************************	\rightarrow	-13								20		Score
E.A.	Moderate/High		Low	Г	High		Low		Low		Wol		LOW		Low			Low		WCI.										WOJ			Low										Hgh			Low								MOT		LOW								** BIBJOOM		Landuse

APPENDIX C1 GENERAL ISSUES TABLE

													•	*	•			•				•	•		•	百
										· · · · · · · · · · · · · · · · · · ·	(public).	harvesting and habitat degradation	Population declines due to over	composition in watershed (Public).	Shifts in natural native fish species	Schell 2002).	(Gottesfeld et al. 2002; Bustard and	Over harvesting of anadromous species	2002)	populations (Public; Gottesfeld et al.	potentially impacting wild fish	Aquaculture/Farming of Atlantic Salmon	High angling pressure (Public)	by sub-species and by watershed (Public)	Genetic diversity of native fish species	FISHERIES MANAGEMENT
0	•	•	• •	Ó									۰	Ó		•		•		• 13		1	•	•	•	
Changes in ocean conditions affecting survival rates of anadromous species.	Global warming / climate change warming water and reducing summer and autumn water flows (public)	Increased another access to sensitive fish habitats (muhlic)	Large volume water utilization associated with proposed industrial projects (Public). Changes in stream and river morphology (public)	Altered wetland integrity and function due to adjacent forest harvesting.	fecundity and survival.	Low flows in summer and fall can lead to migration barriers and reduced fish	Low flows in winter can result in dewatering of redds, freezing.	High flows during migration can, however, improve access to spawning grounds.	habitats.	of inveniles stranding of inveniles or forcing some species to snawn in marginal	and melt, "Ilashy water flows" Gottesfeld et al. 2002; Bustard and Schell 2002).	watersheds (i.e.: groundwater and small stream flow interception, snow accumulation	Timber harvest rates and road building impacts on the hydrological processes of	Water contamination associated with mining activities (Gottesfeld et al. 2002; public).	2002).	Loss of channel stability associated with road stream crossings (Bustard and Schell	associated with forestry and range practices (Bustard and Schell 2002; Public).	Reduced riparian function and integrity, especially along small tributary streams	habitats (Bustard and Schell 2002).	Increased sedimentation affecting water quality, snawning, rearing and overwintering	Beaver dams	 Road crossings 	Barriers to fish migration	Increased water temperatures (Bustard and Schell 2002).	Loss of critical spawning areas in lakes, rivers and streams for all fish species (public).	HABITAT
				•					•		9		•		•		H			•		-	•		•	D)
			established for the Ske	Linkage between escap	changes.	channel form and rates	activity, runoff rates, n	recovery after logging	Information on snowme	quality.	Definition of cumulative	on wetland integrity, fu	Information on the imp	within the watershed.	Baseline data on the cu	riparian management re	temperatures and the in	Definition of the natura	to fish passage.	Number and location o	for concern?	vs. introduced). What	Information on sedime	fish species.	Productive capacity of	DATA GAPS

			ilssies	
Species	Stock Status	Fisheries Management	्रिहोर्गाहा 	Data Gaps
Lake Chub Continued	due to presence in upper Pimpernel Creek above a barrier (P. Giroux, WLAP, in AEM 2001).			
Longnose Suckers	 Present in McBride, Tagit, Owen and Lamprey creeks and associated lakes as well as Nanika-Kidprice lake system (Bustard and Schell	 No species specific issues identified in summary material or public input. 	 No species specific issues identified in summary information. 	distribution (Bustard and Sche
Largescale Suckers	 Common throughout mainstem Morice River (Bustard and Schell 2002). Present in Collins, McBride, Morice and Owen lakes (Applied Ecosystem Management 2001). 	 No species specific issues identified in summary material or public input. 	 No species specific issues identified in summary information. 	Information on stock status, in distribution (Bustard and Sche
Redside Shiners	 Present in Owen, Collins, McBride and Morice lakes and Tagit Creek (Applied Ecosystem Management 2001). 	 No species specific issues identified in summary material or public input. 	 No species specific issues identified in summary information. 	distribution (Bustard and Sche
Prickly Sculpins	 Common in Morice and Nanika rivers and in lower McBride Creek. Also present in Owen and lower Gosnell creeks (Bustard and Schell 2002). 	 No species specific issues identified in summary material or public input. 	 No species specific issues identified in summary information. 	Information on stock status, Indistribution (Bustard and Sche
Pacific Lamprey	 Abundant and widely distributed throughout Morice watershed. Large numbers have been observed spawning in Owen and Lamprey creeks as well as the Morice River (Bustard and Schell 2002). 	 No species specific issues identified in summary material or public input. 	No species specific issues identified in summary information.	distribution (Bustard and Sche

Species	Stock Status	risneries Management	्राज्यात्र ।		-Data Gaps
Pink	Moricetown fishway installed to aid fish	 No species specific issues identified in summary material or 	Sediment inputs and debris loading within diagrams to the	•	Implications of up to 800,000 pink carcass
	migration in 1950.	public input.	mainstem Morice River between	•	Information on specific breakout of the
	 Escapement numbers quite volatile since 1950s 		Gosnell/Thautil confluence and	1	distribution of spawners for monitoring ch
	Morice Pinks make up an		2002).	•	Information on spawning sites (spawner
	average of 9% of the Skeena		 Rain on snow events and fall floods 		densities, incidence of redd crowding and
	pink escapements since		may influence pink spawning		superimposition)
	1985 (Bustard and Schen		locations and winter egg survival		Assessment of the carrying capacity of the
	2002).		partly due to freezing.		Morice River to accommodate pink spawn (Bustard and Schell 2002).
Steelhead	 Bulkley-Morice sub-unit is 	 High Alaskan and Canadian 	 Sediment inputs and debris loading 	•	Need a solid juvenile and adult database for
	the most important	commercial bycatch (Bustard and	within direct tributaries to the		management of this stock.
	steelhead system in the	Schell 2002)	mainstem Morice River between	•	Information on bottlenecks to smolt produc
	1999-2000 mark-recapture	(minor hooking mortalities	Owen Creek	•	Information on adequate spawner recruitm
	estimates indicate 4000 to	(Hooton) in Bustard and Schell	 Sediment inputs into Owen, Fenton, 	•	tributaries and the mainstem Morice River
	Morice	• Native fishery rates (Rustard and	Peacock and Lamprey watersheds		subsequent downstream movements from I
	 Catch and release only 	Schell 2002)	particularly in Lamprey and Owen		ाउँ आगि। अबहुट (जिन्नाच बाच उटाहा। 2002).
	regulation in effect for sport	 Introduction of escaped Atlantic 	creeks (Bustard and Schell 2002).		
	the 1000's	salmon to the Skeena may threaten	 Low summer flows restricting 		
	Wet'suwet'en food fishery	2002, public)	especially in Lamprey and Owen	ii H	
	records at Moricetown		creeks (Bustard and Schell 2002)	Ī	
	fish caught annually since			-	
	the 1980's (Bustard and				
	Schell 2002).				
	throughout the Morice	angling regulations considering the	habitat areas (Bustard and Schell	•	size at age data
19	watershed (Morice, Owen,	unproductive stocks.	2002).	•	Information on and location of spawning at
1	Lamprey, Phipps and Bill Nye lakes, Nanika-Kidnrice		Dewatering prior to fall emergence	•	Genetic information on juvenile rainbow in
	lakes system, Houston		(Bustard and Schell 2002)		they are progeny of steelhead or resident
	Tommy, Fenton, and				rainbow (Bustard and Schell 2002).
	River.)				

Morice Watershed-based Fish Sustainability Plan - Stage II

	I LOGO SU COLUMNICA III SICCE			The state of the s
	Dood stream crossings in steen		Relatively low densities in	
(Bustard and Schell 2002).	forest roads		streams	
Distribution is limited to point	small seepage flows by adjacent		catches in many smaller	神経 神子に という
migration patterns and home ra	 Interception of ground water and 	public input.	watershed and dominate the	
overwintering locations, popul	temperatures.	identified in summary material or	species in the Morice	
	 Riparian management and instream 	 No species specific issues 	Most widely distributed	Dolly Varden
	(Bustard and Schell 2002; public).			
	areas and other key habitats			
	 Improved angler access to staging 		species.	
	2002).		Provincially blue-listed	
	forest roads (Bustard and Schell		Schell 2002).	
	small seepage flows by adjacent		individuals (Bustard and	
key rearing areas (Bustard and	Interception of ground water and	access.	population < 1000	
Information on location and ch	low water temperatures	unnroductive stocks and anoler	Possible snawning	
watershed carrying capacity.	concern because built front require	Sport fishery narvest rates - fiberal angling regulations considering the	Slock status is currently	• morr mg
125	IIi		and Schell 2002).	
			lakes and ponds (Bustard	
			Holland and many smaller	
			Collins, Chisholm, Julian	
			within watershed (McBride,	
			associated with tributaries	H
			Present in larger lakes	
			the Gosnell watershed.	
	habitats (Bustard and Schell 2002).		Widespread distribution in	
relationships (Bustard and Schell 2)	 Increased angler access to key 	public input.	and Nanika River	
migration patterns, stock status and	stream crossings.	identified in summary material or	Morice River, Morice Lake	
Information on spawning, overwing	Fish passage associated with road	 No species specific issues 	Uncommon in the mainstem	Cutthroat I rout
			and Schell 2002).	
			1980 to present (Bustard	
			spawning period imposed in	
			fishery during Chinook	
			upper Morice River sport	
	ń.		Regulations closing the	
			Morice watersned.	
		The second secon	Mario Wetarched	
		The second secon		Continued
2000			Adults occur in low	Rainbow Trout
	<u>"labirai</u>	Fisheries Management	Stock Status	Species
A 10 COLUMN 18 18 18 18 18 18 18 18 18 18 18 18 18				

Phononical Contract of the Con

	Ž
	걸.
	Norice \
	_
	ate
	Vaters
	he
	ä.
	ba
	Se
	atershed-based Fi
	Fish
	7
	Ĕ
á	ळ
	na
	₽.
	lity Pla
	D
	3
	1
	St.
	tage
	age II

			SSIES	
Species	Stock Status	Fisheries Management	Habitat .	Data Gaps
bolly Varden Continued	most stream systems Stocks not threatened or declining in Morice watershed (Bustard and Schell 2002).		gradient streams (Bustard and Schell 2002).	
19ke frout	 Present in four lakes (Morice, Owen, McBride and Atna) (WLAP Lake Surveys; Bustard and Schell 2002). 	 Sport fishery harvest rates - liberal angling regulations considering the unproductive stocks. 	 No species specific issues identified in summary information. 	 Information on stock status, migratory cand spawning areas (Bustard and Schell)
Kokance	 Only reported in two lakes (Morice and Shea) (Bustard and Schell 2002). 	 No species specific issues identified in summary material or public input. 	 No species specific issues identified in summary information. 	 Information on stock status, spawning and distribution (Bustard and Schell 2002).
Whitefish	 Three species of whitefish reported in Morice watershed (Applied Ecosystem Management 2001). 	 No species specific issues identified in summary material or public input. 	 No species specific issues identified in summary information. 	 Information on stock status, spawning name and distribution (Bustard and Schell 200
	 Pygmy whitefish in Owen and Morice lakes, lake whitefish in McBride and Morice lakes and mountain whitefish spread throughout the watershed 			
	 Mountain whitefish are the most abundant species throughout mainstem rivers (Bustard and Schell 2002). 			
Burbot	 Present in McBride, Morice and Owen lakes (Applied Ecosystem Management 2001). 	 No species specific issues identified in summary material or public input. 	species specific issues identified ummary information.	Information on stock status, life history
Lake Chub	 Present in Gosnell, Thautil, Lamprey and Owen creeks as well as Morice and Owen lakes (Applied Ecosystem Management 2001). Regionally important status 	No species specific issues identified in summary material or public input.	• No species specific issues identified in summary information.	distribution (Bustard and Schell 2002).

APPENDIX C2 SPECIES SPECIFIC ISSUES

		· · · · · · · · · · · · · · · · · · ·	Issues	
Snecies	Stock Status	Fisheries Management	Habitat	Data Gaps
		First Nations in-river harvest rate	Naturally low productivity in Morice	Beach spawning sites in Morice a
Sockeye	• Spawner recruitment generally low since the 1950's (Bustard and Schell	(Bustard and Schell 2002) Mixed fishery bycatch (Public).	, ,	
	Hatchery built (pilot project) to increase stocks –		 Sediment inputs to spawning grounds from Glacier Creek (Bustard and Schell 2002). 	and subsequent collapse of socke escapements (Bustard and Schell
Chinook	unsuccessful.	First Nations, Commercial and	Sediment inputs and debris loading	Information on spawner abundan
CALL	Stocks Stock escapement is stable (Bustard and Schell 2002).	Recreational harvest rates (Bustard and Schell 2002).	within direct tributaries to the mainstem Morice River especially between Gosnell/Thautil confluence and Owen Creek (Bustard and Schell	Analysis of Morice watershed ca for juvenile rearing (Bustard and
			 2002). Redd superimposition (a key limiting factor). 	
Coho	Conservation management strategies implemented in 1998	 Mixed fishery bycatch (public and technical committee) High Alaskan fishery harvest rate 	 Sediment inputs and debris loading within direct tributaries to the mainstem Morice River especially 	 Information on physical characters spawning habitats. Information on habitat capacity
History Area Area Area Area Area Area Area Area	Strategic stock enhancement conducted 1998-1999	(Bustard and Schell 2002)	and Owen Creek (Bustard and Schell	for juveniles Stock identification (Bustard and
	Morice populations recovering since 1998		 Poor ocean survival (Bustard and Schell 2002) 	OLOGA ROSESSA
	Decreasing cono composition in juvenile comples (Bustard and Schell		 Access to spawning sites due to low fall flows and beaver dams especially 	
	2002).		in Owen, Lamprey and McBride creeks (Bustard and Schell 2002).	
uk o			dissolved oxygen in small rearing streams (Bustard and Schell 2002).	
er (1000) Karthari Tankari Mariana			 Dewatering of rearing and overwintering habitats (Bustard and 	
	5 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		Fish passage associated with road ctream crossings (Public).	

APPENDIX D POSSIBLE SCORING MATRIX TO RANK ISSUES

Potential Impact of Issue			Ra	ting	
	VH	Н	M	L	Score
Risk (multiply a x b)					生物 经保险
a) probability of occurrence					
b) magnitude of impact	+				1
Real or perceived risk(4 = real; 1 = perceived)		•			A Philippins
Seriousness		T			
a) Direct impact to a stock at risk					America
b) Number of species directly benefiting by addressing					THE PROPERTY OF
issue				1	
Urgency					Supple And F
Subtotal		KEN TO	4	ALC: N	LAUNA CONT.

Ability to Plan to Address Issue			Ra	ting	
	VH	Н	M	L	Score
Feasibility / implementer's ability to influence change					944994
Willingness of impacter to change					September 1981
Time needed to deal with issue (long = Low, short = VH)					STATISTICS.
Complexity of issue					HARRY CO.
Subtotal	dra Surface	14 4 年	di ten	State of	TO SERVICE
TOTAL for issue	Shrop-art	#W. 4	A Albah	ARMINAL N	MANAGEMENT .

Directions

Place scores in shaded boxes

VH =	Very High	4 points
H =	High	3 points
M =	Moderate	2 points
L=	Low	1 point

Definitions:

Risk: the probability of occurrence X magnitude of impact of an issue

Real or Perceived Risk: perceived risk is risk for which there is only intuitive or anecdotal evidence. Real Risk is based on empirical or scientific evidence.

<u>Seriousness:</u> Is the issue acute or dangerously chronic? Is the issue widespread or highly localized?

Urgency: How quickly does action need to be taken?

<u>Feasibility / ability to influence change:</u> What is the implementer's ability to influence changes needed or to overcome challenges associated with the issue.

<u>Willingness of impacter to change</u>: are those responsible for management willing to change their practices?

<u>Time needed to deal with issue</u> (long = Low, short = VH): Amount of time and resources needed to deal with the issues relative to other issues.

Complexity of issue: How multifaceted is the issue? Do we understand the issue – the cause and effect? Does the issue deal with a large number of stakeholders?

Addendum to:

A Plan to Conserve and Protect Morice Watershed Fish Populations and their Habitat – Stage II (Tamblyn and Croft, 2003)

Following the distribution of the final draft of *A Plan to Conserve and Protect Morice Watershed Fish Populations and their Habitat – Stage II* (Tamblyn and Croft, September 2003), Community Futures Development Corporation of Nadina received feedback we felt was important to pass to people receiving copies of the report.

Concerns were expressed over the wording of the following statement found in **Section 5.2.1 Priority Fisheries Management Issues** on page 24:

"Past high First Nation in-river harvest rates for sockeye and steelhead during some years"

Comments received:

- This statement should not be considered a priority or significant fishery issue.
- First Nations have a traditional right to harvest fish.
- The statement may lead to misconceptions by certain members of the public about the numbers of fish caught by the Wet'suwet'en.
- There is no current conservation concern regarding Bulkley steelhead populations.

We would like to point out that Wet'suwet'en Fisheries have been leaders in local stock assessment efforts over the past several years. They have run, or been partners in, projects to improve fisheries management in the Bulkley and Morice watersheds including coho, steelhead and sockeye mark-recapture programs. Currently, the Wet'suwet'en have voluntarily stopped harvesting sockeye at Moricetown in order to monitor stocks and allow the Maxan and Nanika sockeye populations to rebuild.

The Wet'suwet'en have also been a valuable member of the Bulkley Morice Salmonid Preservation Group and have participated in Strategic Planning for this group and are members of both the Morice WFSP planning and technical committees.

Sincerely,

Greg Tamblyn WFSP Planner

They Lambly