

THE BABINE LAKE SALMON DEVELOPMENT PROGRAM

DFO - Library / MPO - Bibliothèque



12001682

PROGRESS REPORT
TO MARCH 31, 1968

PREPARED JOINTLY BY:

THE DEPARTMENT OF FISHERIES
OF CANADA

THE FISHERIES RESEARCH BOARD
OF CANADA

Report 1968-7

AUGUST, 1968
VANCOUVER, B. C.

SH
157.85
.S6
B32
D.7

LIBRARY
FISHERIES AND OCEANS
BIBLIOTHÈQUE
PÊCHES ET OCÉANS

70092

SH
157.85.56
B32
D

THE BABINE LAKE SALMON DEVELOPMENT PROGRAM

PROGRESS REPORT TO MARCH 31, 1968.

THE LIBRARY
BEDFORD INSTITUTE OF
OCEANOGRAPHY
BOX 1006
DARTMOUTH, N.S. B2Y 4A2

Prepared jointly by the Department of Fisheries
and the Fisheries Research Board of Canada.

Vancouver, B. C.

TABLE OF CONTENTS

	Page
I. INTRODUCTION	1
II. SUMMARY OF ENGINEERING PROGRAM	
A. Introduction	3
B. Description of Facilities	
1. Fulton River	
(a) Project No. 1 - Spawning Channel No. 1	4
(b) Project Nos. 3 and 4 - Tunnels and Regulating Works	4
(c) Project Nos. 9 and 16 - Spawning Channel No. 2	7
(d) Project No. 15 - Pipeline	7
(e) Project No. 11 - Fulton Storage Dam	8
(f) Project Nos. 6, 7, 8, 12, 13 and 14 - Reservoir Clearing	8
2. Pinkut Creek	
(a) Project No. 2 - Taltapin Lake Control Works	9
(b) Project No. 5 - Pinkut Creek Spawning Channel	9
C. Construction Program Time-Table and Yearly Expenditures	10
D. Other Studies	
1. Hydrometeorological Program	10
2. Morrison River	13
III. SUMMARY OF BIOLOGICAL PROGRAM	
A. Introduction	14
B. Comparison of Fry Production from Natural and Artificial Spawning Grounds	
1. Fulton River	
(a) Natural Spawning Grounds	15
(b) Fulton Spawning Channel No. 1	17
2. Pinkut Creek (15 Mile Creek)	19

TABLE OF CONTENTS (continued)

	Page
III. SUMMARY OF BIOLOGICAL PROGRAM (continued)	
C. Comparison of Fry Produced Naturally with those Produced using Fish Cultural Aids	
1. Studies of Behaviour, Physiology, and Performance	
(a) Fry Migration Timing	21
(b) Fry Quality Studies	23
(i) Length, weight and development stage	23
(ii) Swimming performance	23
(iii) Starvation resistance	24
(iv) Vulnerability to predation	25
(v) Biochemical composition	26
2. The Distribution, Growth, and Survival of Sockeye Fry Produced from the Fulton River and Artificial Spawning Channel in 1966	27
D. Primary Productivity of Babine Lake	33
E. Other Studies	
1. Adult and Smolt Sockeye Enumerations	34
2. Factors Associated with Age at Maturity of Skeena River Sockeye	42
3. A Review of the Literature on the Effects of Changes in Temperature Regime on Developing Sockeye Salmon Eggs and Alevins	
(a) Introduction	44
(b) Effects During Period Before Egg Deposition	47
(c) Effects During the Larval Stages	48
4. Diel Vertical Movement of Pelagial Sockeye Salmon Juveniles	
(a) Problem and Methods	62
(b) Early Results	63
IV. ACKNOWLEDGEMENTS	71
V. LITERATURE CITED	72

*Manu. 949.
FRBe.*

LIST OF FIGURES

	Page
1. Babine Lake, British Columbia.	2
2. Fulton River General Layout	5
3. Sockeye fry migration, Fulton River and Channel, 1967.	22
4. Locations of marked sockeye recoveries in Babine Lake, 1966.	29
5. Estimates of the relative density and abundance of marked and unmarked sockeye by lake area and fishing period. See Figure 4 for location of areas.	30
6. Babine Lake sockeye smolt production.	37
7. Age composition - Babine sockeye.	39
8. Parasitism of Babine sockeye smolts by tapeworm <u>Eubothrium salvelini</u> in 1966 and 1967.	43
9. Plot of midwater trawl catch per minute of juvenile sockeye and the simultaneous number of echo sounder targets per minute in the 10-ft. stratum of water at the depth of the trawl. Data from August, 1967 in the North Arm of Babine Lake.	65
10. Echo sounder targets per minute (.3 mile) by 10-ft. strata during the 24-hour series of August 5-6 in the North Arm.	66
11. Echo sounder targets per minute (.3 mile) by 10-ft. strata during the 24-hour series of October 2-3 in the North Arm.	68
12. The diel vertical distribution of zooplankton species in the North Arm of Babine Lake on September 22-23, 1967. Each depth is expressed as a percent of the number of that species taken in a particular vertical profile.	69

LIST OF TABLES

	Page
I. BABINE LAKE DEVELOPMENT PROGRAM CONSTRUCTION SCHEDULE	11
II. BABINE LAKE DEVELOPMENT PROGRAM REVISED COST ESTIMATES	12
III. SUMMARY OF FULTON RIVER EVALUATION PROGRAMS, 1961 TO 1967	16
IV. SUMMARY OF FULTON SPAWNING CHANNEL EVALUATION PROGRAMS, 1965 TO 1967	18
V. SUMMARY OF PINKUT CREEK EVALUATION PROGRAMS, 1963 TO 1967	20
VI. CHI-SQUARE TESTS OF DEPARTURES FROM EXPECTED RATIO OF RIVER AND CHANNEL RECOVERIES, 1966	32
VII. BABINE SOCKEYE ESCAPEMENTS, IN THOUSANDS OF FISH, 1949-1966	36
VIII. AGE COMPOSITION OF SKEENA STOCKS OUTSIDE BABINE, 1963-1967	41
IX. EXPECTED RELATIONSHIP BETWEEN DEVELOPMENT RATE, FERTILIZATION TO HATCHING, AND TEMPERATURE	52
X. DEVELOPMENT RATES AT DIFFERENT TEMPERATURES	53

I. INTRODUCTION

Babine Lake (Figure 1), supports one of the largest sockeye salmon runs in British Columbia. It produces the bulk of the Skeena River catch, which currently averages about 500,000 sockeye annually and which has a landed value in excess of one million dollars.

The lake is comprised of two major areas. The North Arm contains about 12% of the total lake surface area, and the main lake contains the remainder. Spawners are distributed nearly equally between the Babine River spawning grounds, adjacent to the North Arm, and the streams tributary to the main lake.

Studies conducted by the Fisheries Research Board of Canada have indicated that sockeye fry densities are consistently lower in the main basin than in North Arm, despite similar densities of planktonic organisms. Basic productivity, as measured by the carbon-14 in situ method, is at least as high, on a per unit area basis, in the main basin as in the North Arm (Narver, 1967). This evidence suggests that the lower sockeye density in the main basin is due not to a limited food supply but to limited fry production by its tributary streams. In short, the streams tributary to the main basin are considered incapable, in their natural state, of producing enough sockeye fry to fully utilize the lake's potential as a nursery area.




Figure 1. Babine Lake, British Columbia

Realizing the development possibility of such a situation, the Resource Development Branch of the Department of Fisheries of Canada embarked, in 1965, on a major program of artificial spawning channel construction coupled with flow control to increase fry production in spawning streams tributary to the main basin. The streams selected were Fulton River and Pinkut Creek. This report deals with the progress which has been made in the construction and evaluation of the Babine Lake Sockeye Salmon Development Program up to March 31, 1968.

II. SUMMARY OF ENGINEERING PROGRAM

A. Introduction

Following approval of the basic development project early in 1965, detailed engineering studies were initiated and the first spawning channel was completed at Fulton River in October, 1965. Design work on the overall project has continued, and definite plans for the storage dams, control works, water conduits, and spawning channels have been formulated. The design work has now been completed, with the exception of the second spawning channel at Fulton River.

The estimated cost for the complete program, based on construction costs at May, 1967, is 7.8 million dollars. This represents an increase of 2.5 million dollars to the original estimate prepared in 1964. This increase is attributable to: (1) the significant increase in



Fulton River falls and valve house
under construction at the outlet of
the regulatory tunnel

construction costs which have occurred since 1964; (2) the higher than anticipated construction costs in the Babine area itself; and (3) the design of more complex and therefore more expensive structures than had been envisaged on the basis of preliminary data. The estimated benefit cost ratio based on the increased construction costs for the total program exceeds 3:1.

B. Description of Facilities

1. Fulton River

(a) Project No. 1 - Spawning Channel No. 1

Spawning Channel No. 1, the first of the development projects, was completed in October, 1965 at a total cost of \$500 thousand.

The channel (Figure 2 and cover) is 4,900 feet long including pools and has a bottom width of 30 feet and a gradient of 9 feet in 10,000 feet. At a discharge of 75 cfs the depth of water in the channel is 1.3 feet and the average velocity 1.8 feet per second. The channel has an estimated capacity for 24,000 adult sockeye in 15,000 square yards of spawning gravel.

(b) Project Nos. 3 and 4 - Tunnels and Regulating Works

The regulating works are shown schematically in Figure 2. Provision has been made in the height of the gate structure for an increased reservoir elevation if additional flow control is required in the future.

The gate system has been so arranged that water may be drawn from several different depths in the reservoir to obtain the required temperatures. If the water in the Fulton Lake outlet basin is not sufficiently cool, a pipe will be placed in the intake channel and extended out to a deeper area of the lake. Provisions have been made in the regulating structure to accommodate a low level intake gate (see Figure 2).

Regulated flows to Fulton River will range from 100 cfs during the spawning and incubation period to 3500 cfs during spring freshets. Outflows will be passed through a concrete lined tunnel and regulated by two 84 inch and one 30 inch diameter hollow cone valves.

Flows to spawning channel no. 2 will range from 150 cfs during the spawning period to 100 cfs during the incubation period. Water will be conveyed through a 7.5 foot modified horseshoe-shaped tunnel 3,850 feet in length.

Project No. 3, which included rock excavation for the tunnels and the regulating and control structures, was completed in 1966 at a cost of \$585,000. Although original studies had suggested that the supply tunnels would not require lining, a detailed study of rock conditions after excavation resulted in plans to line them with concrete.

Project No. 4, which included construction of the regulating works and lining of the tunnels, began in July 1967 and is scheduled for completion in May 1968 at a cost of \$945,000. The cost of the supply tunnel lining represents approximately 25% of this total cost. In addition, seven supply contracts for equipment and metalwork were awarded at a total cost of \$300,000.

(c) Project Nos. 9 and 16 - Spawning Channel No. 2

Spawning Channel No. 2 will be 17,000 feet long, with a bottom width of 50 feet and a gradient of 20 feet in 10,000 feet. At a discharge of 150 cfs the depth of water in the channel will be 1.25 feet and the average velocity 2.2 feet per second. The channel will accommodate an estimated 151,000 adult sockeye and will produce 78 million fry at an egg-to-fry survival rate of 40%.

The channel will be constructed in two stages. The first stage is scheduled for completion in August 1969 and the second stage in 1970. The estimated cost is 1.8 million dollars.

(d) Project No. 15 - Pipeline

A 6,000 foot long pipeline is required to carry water from the supply tunnel to spawning channel no. 2. The project will commence in 1968 and be completed in 1969 at an estimated total cost of \$525,000.

(e) Project No. 11 - Fulton Storage Dam

The storage dam at the outlet of Fulton Lake will be a concrete gravity structure having a spillway crest approximately 25 feet above the low water level of the lake. It will impound 76,000 acre feet of water. Two outlet gates will be placed in the dam and will have a minimum capacity of 100 cfs. These gates will permit water to be drawn from the surface of the reservoir and released into the Fulton River during the spawning period.

Construction of the dam is scheduled to commence in August 1968 and to be completed by April 1969, in time to store water from the spring runoff. The estimated cost of the project is \$600,000.

(f) Project Nos. 6, 7, 8, 12, 13 and 14 - Reservoir Clearing

A comprehensive field investigation was carried out in 1967 and a program for reservoir clearing was formulated to allow storage of water by the spring of 1969. Five major contracts will be awarded in 1968 for logging and slashing. The resultant slash will be piled and burned. Log booms will be constructed and positioned in Fulton Lake to collect all floatage. In 1969, all floatage will be collected, sorted and burned. The lake and shoreline will be swept and cleaned in 1970. The

estimated total cost for the reservoir clearing is \$750,000.

2. Pinkut Creek


(a) Project No. 2 - Taltapin Lake Control Works

A dam and outlet control works was constructed at Taltapin Lake in 1966 at a cost of \$200,000. It has 6 1/2 feet of live storage with a reservoir capacity of 32,000 acre feet and will provide a minimum discharge from Taltapin Lake of 75 cfs during the spawning and incubation periods. The length of the spillway in the dam was designed to maintain the normal spring runoff within the present maximum lake levels. This feature was added after consultation with the B. C. Forest Service and Department of Recreation and Conservation, and will ensure minimum effects on the lakeshore area.

In order to realize the total storage of 6 1/2 feet it will be necessary to dredge the area immediately below the outlet of Taltapin Lake. This work is scheduled for 1968, at a cost of \$50,000.

(b) Project No. 5 - Pinkut Creek Spawning Channel

Water will be diverted from Pinkut Creek to the proposed spawning channel by a small diversion dam in the canyon area, near the upper limits of the present spawning grounds. A 750 foot long tunnel will convey water to the channel.



The Fulton River Regulatory Works showing the valve house and two - 84 inch discharge regulators

The spawning channel will be completed in two stages. The first stage, to be completed in 1968, will have an overall length of 10,300 feet, a bottom width of 40 feet and a gradient of 9 feet in 10,000 feet. At a discharge of 75 cfs, the depth of water in the channel will be approximately 1.25 feet and the average velocity 1.8 feet per second. The channel will provide spawning area for 67,000 adult sockeye and produce 34 million fry at an estimated egg-to-fry survival rate of 40%. The second stage will be of equal size but will be deferred until utilization of the first stage has been achieved.

C. Construction Program Time-Table and Yearly Expenditures

The present construction schedule is outlined in Table I. Numerous changes were made in the original schedule as funds became available and as requirements and designs were modified and finalized.

The present Babine Lake Development Program will cost 7.8 million dollars, a sizeable increase over the original estimate of 5.28 million approved in principle by Treasury Board in 1965. Yearly expenditures are detailed in Table II.

D. Other Studies

1. Hydrometeorological Program

The Babine Lake hydrometeorological program co-ordinates the collection and treatment of all

TABLE I. BABINE LAKE DEVELOPMENT PROGRAM
CONSTRUCTION SCHEDULE

LOCATION	1965-66	1966-67	1967-68	1968-69	1969-70	1970-71
FULTON RIVER	- <u>Project No. 1</u> Spawning Channel No. 1 (Completed Oct. 15/65)	- <u>Project No. 3</u> Diversion & Water Supply Tunnels	- <u>Project No. 4</u> Regulating Works, Diversion & Water Supply Tunnel Linings	- <u>Projects 6, 7, 8, 12, 13</u> Reservoir Clearing	- <u>Project No. 14</u> Reservoir Clearing	- <u>Project No. 16</u> Spawning Channel No. 2, Stage 2
		- Bridge across Spawning Channel at Highway	- Relocation of Strimbold Mill	- <u>Project No. 9</u> Spawning Channel No. 2, Stage 1	- <u>Project No. 9</u> Completion of Spawning Channel No. 2, Stage 1	- <u>Project No. 18</u> Reservoir Clearing
				- <u>Project No. 11</u> Fulton Lake Storage Dam	- <u>Project No. 11</u> Completion of Fulton Lake Storage Dam	
				- <u>Project No. 15</u> Pipeline to Spawning Channel	- <u>Project No. 15</u> Completion of Pipeline	
PINKUT CREEK		- <u>Project No. 2</u> Taltapin Lake Control Works	- Spawning Channel Stage 1	- <u>Project No. 5</u> Completion of Spawning Channel Stage 1		
		- Access Road to Spawning Channel Site	- Completion of Access Road Surfacing	- <u>Project No. 10</u> Excavation down- stream of Taltapin Lake Control Works		
	\$500,000	\$1,100,00	\$1,400,000	\$2,355,000	\$1,600,000	\$810,000

TABLE II. BABINE LAKE DEVELOPMENT PROGRAM -
REVISED COST ESTIMATES

	Estimated Cost 1964	Actual or Estimated Cost 1967
I. <u>FULTON RIVER</u>		
1. Spawning Channel No. 1 Completed in 1965	\$ 425,000	\$ 500,000
2. Dam, Reservoir Clearing, and Mill Relocation	950,000	1,500,000
3. Diversion Tunnel and Control Works	200,000	1,100,000
4. Spawning Channel No. 2 Including Water Supply Tunnel and Pipeline	2,200,000	3,200,000
MISCELLANEOUS		350,000
II. <u>PINKUT CREEK</u>		
5. Storage Dam and Channel Dredging	200,000	250,000
6. Access Roads	80,000	100,000
7. Spawning Channel	1,225,000	800,000
TOTALS	<u>\$5,280,000</u>	<u>\$7,800,000</u>

Capital 7,347,000 to
end of 1969
Operation \$ 320,000
permanent salaries &

hydrologic and meteorological data in the area to provide basic information for the design and operation of spawning channels and the regulation of reservoirs.

Of the several studies being undertaken, the most important is attempting to determine the effects of climatological factors, especially solar radiation, on spawning channels in late August and September. It is possible that warming during the day will produce adverse water temperatures in long spawning channels unless proper protection is designed for them.

Fulton River spawning channel no. 1 is being used as a test channel and the results will be applied to the new channels.

Other work is presently being undertaken with respect to: (1) Fulton Lake reservoir water supply temperature control; (2) Fulton River watershed spring runoff forecasting; (3) stream runoff records at Fulton and Morrison Rivers and Pinkut Creek; and (4) basic climatological data collection.

2. Morrison River

Preliminary field surveys have been conducted at Morrison River by the Department of Fisheries of Canada. Examination of the topography and water surface profiles suggests that an initial spawning channel development in the upper river area would provide 30,000 square yards of gravel and accommodate

43,600 adult sockeye. The channel could eventually be extended to the lower river area to provide an additional 49,000 square yards of spawning area.

III. SUMMARY OF BIOLOGICAL PROGRAM

A. Introduction

The present biological program is proceeding upon the following premises: (1) that the main basin of Babine Lake is presently underutilized and can support more sockeye fry if these were to be produced; (2) that these large numbers of additional sockeye fry can be produced in artificial spawning channels and in natural channels with regulated flow; and (3) that the fry so produced are entirely comparable to naturally produced fry in their ability to survive to the adult stage.

There is considerable evidence supporting the above premises, reviewed in the report "Proposed Sockeye Development Program for Babine Lake". It is one of the objectives of the evaluation program, however, to continually assess the validity of these assumptions. This type of evaluation program is the only rational and objective basis for the improvement of enhancement techniques and ultimately for obtaining the maximum utilization of the potential of Babine Lake.

B. Comparison of Fry Production from Natural and Artificial Spawning Grounds

1. Fulton River

(a) Natural Spawning Grounds

The data collected at Fulton River is summarized in Table III. Actual counts of adult spawners have been made since 1962, prior to which the tag and recovery method was utilized. The adults were sampled for sex and age composition, fecundity, and retention. Fry production was measured with vertical samplers until converging throat traps, mounted on a fence, were used to assess the 1966 brood year (Walker et al, in press). The latter method is the more accurate of the two and will be used in all future Fulton River downstream enumerations.

The data indicated an inverse correlation between actual deposition and percent survival, although this has not been shown to be statistically significant. If a correlation does exist it is probably the result of superimposition.

It is important to note one serious limitation of the data. There has undoubtedly been considerable error in enumeration, particularly during periods of high water. During the 1966 program, for example, debris carried downstream by the high spring runoff prevented continuous operation of the

TABLE III. SUMMARY OF FULTON RIVER EVALUATION PROGRAMS, 1961 TO 1967

	Brood Year							1968
	1961	1962	1963	1964	1965	1966 ¹	1967 ¹	
Migration Pattern								
5% of Run	-	Aug. 23	Aug. 22	Aug. 25	Aug. 27	Aug. 15	Aug. 27	
50% of Run	-	Sept. 6	Sept. 7	Sept. 6	Sept. 7	Sept. 7	Sept. 10	
95% of Run	-	Sept. 21	Sept. 19	Sept. 23	Sept. 20	Sept. 19	Sept. 21	
Number of Spawners	189,000	86,000	154,000	120,000	139,000	62,000	118,000	100,000
Sex Composition								
% Females	45.2	52.0	35.2	52.0	52.3	38.1	47.5	
% Males	44.6	48.0	29.2	45.3	36.4	27.0	49.3	
% Jacks	10.2		35.6	2.7	11.3	34.9	3.2	
Number of Females	85,428	44,720	54,208	62,400	72,763	23,071	56,284	16
No. Eggs/Female	2,887	3,272	3,013	3,103	2,910	3,204	3,099	190.0
Potential Deposition (millions)	246.8	146.3	163.3	193.6	211.7	73.9	174.4	
% Retention	3.7	6.7	9.4	3.3	14.1	1.2	1.6	
Apparent Deposition (millions)	237.7	136.5	148.0	187.0	189.0	73.0	171.6	
Fry Production (millions)	26.5	41.7	46.5	24.5	-	21.7	-	32.0
% Survival	11.0	30.5	31.4	12.5	-	29.7	-	15.0
Fry Weight (grams)	-	.19	.19	.18	.15	.16	-	
Fry Length (millimeters)	-	31.3	30.3	28.2	29.5	29.4	-	
Fry Migration								
Start	May 1	April 25	April 30	April 28	April 30	May 1	-	
Peak	May 28	May 23	June 4	-	June 6	June 2	-	
End	June 21	June 12	June 17	-	June 16	June 15	-	

¹The 1966 and 1967 figures exclude those sockeye which utilized the spawning channel (see Table IV).

sampling gear, and the peak of the fry migration was not measured (note absence of 1965 fry figures in Table III). The new fence and trapping facilities have eliminated this problem and more accurate population estimates will be obtained in future years.


(b) Fulton Spawning Channel No. 1

The operation to date of Fulton spawning channel no. 1 is summarized in Table IV.

The channel was completed in 1965, but too late for the entry of the natural spawning population of sockeye. Therefore, 1.2 million sockeye eggs were collected from Fulton River spawners, fertilized, incubated to the eyed egg stage, and placed in the channel by the end of November. An 85% survival was obtained from this egg plant.

Adult sockeye were allowed to enter the spawning channel in 1966 (28,000) and 1967 (22,471). The 1966 spawning produced 25.5 million fry, representing a survival of 69%. Survival of the 1967 brood year is indicated by winter sampling to be about 65% up to the end of March.

A winter program is presently being carried out in the spawning channel in an attempt to determine the timing and causes of sub-gravel



The combination adult and fry enumeration weir at Fulton River.
The downstream migrant traps are shown raised out of the water.

TABLE IV. SUMMARY OF FULTON SPAWNING CHANNEL
EVALUATION PROGRAMS, 1965 TO 1967

	Brood Year			
	1965	1966	1967	1968
Number of Spawners	-	28,000	22,471	28,000
% Females	-	41.6	47.5	
Number of Females	-	11,600	10,696	
Eggs/Female	-	3,204	3,099	
Potential Deposition (millions)	-	37.17	33.15	59.0
% Retention	-	1.09	0.99	
Apparent Deposition (millions)	1.20	36.76	32.75	
Number of Fry (millions)	1.02	25.5	16.0	24.7
% Survival	85.0	69.4	49.0	44.7
Fry Weight (grams)	.15	.15	-	
Fry Length (millimeters)	28.8	29.3	-	

mortality. An attempt is being made to correlate egg and alevin mortality with such environmental factors as sub-gravel velocity, stream discharge, water temperature, sub-gravel dissolved oxygen levels, siltation, and ice cover.

Results to date indicate that the heaviest mortality occurs at the time of spawning (cf. 69 and 65% survivals from spawning and 85% survival from planting) while a small additional mortality (approximately 5%) occurs within the first two months of incubation but cannot as yet be ascribed to any one environmental factor.

2. Pinkut Creek (15 Mile Creek)

Biological data has been collected at Pinkut Creek since 1963 and is summarized in Table V. The 1963 population was estimated by the Conservation and Protection Branch but subsequent population figures are tower counts made by the Resource Development Branch.

All fry population estimates for Pinkut Creek were obtained by hydraulic sampling, and are subject to all the limitations of this method. Downstream fry enumeration programs were attempted in 1966 and 1967 to provide a check on the hydraulic sampling results. Unfortunately the downstream sampling was ineffective due to a combination of high river discharges, debris and rising lake levels.

TABLE V. SUMMARY OF PINKUT CREEK EVALUATION PROGRAMS, 1963 TO 1967

	Brood Year					1968
	1963	1964	1965	1966	1967	
Number of Spawners	65,000	146,000	34,000	28,428	33,413	10133*
Migration Pattern						
5% of Run	-	Aug. 25	Aug. 26	Aug. 28	Aug. 25	-
50% of Run	-	Sept. 9	Sept. 4	Sept. 8	Sept. 9	20
95% of Run	-	Sept. 24	Sept. 18	Sept. 18	Sept. 18	1
Sex Composition						
% Females	39.0	56.0	53.0	30.0	42.0	
% Males	61.0	43.0	17.0	45.5	53.0	
% Jacks		1.0	30.0	24.5	5.0	
Number of Females	20,500	76,000	18,000	8,528	14,033	
Eggs/Female	2,930	2,550	3,070	2,940	3,162	
Potential Deposition (Millions)	60.0	228.0	55.0	27.0	44.4	17.8
% Retention	4.0	1.0	7.0	0.4	7.1	
Actual Deposition (Millions)	57.6	226.0	51.2	26.9	41.2	
Fry Production (Millions)	11.0	4.5	6.9	3.7	-2.0	1,883
% Survival	19.0	2.0	13.5	12.5	*	9.8

Spawning channel spawners.
 egg dep.

14,501
 30.2

* Stream only

There has been no consistent correlation between number of females and egg-to-fry survival, and therefore no evidence for mortality due to redd superimposition. This conclusion must be a tentative one, however, because of the limitations of the data.

When the Pinkut Creek spawning channel is completed a full winter program, similar to that at Fulton River, will be carried out.

C. Comparison of Fry Produced Naturally with those Produced using Fish Cultural Aids

1. Studies of Behaviour, Physiology, and Performance

(a) Fry Migration Timing

Truly comparable data is available only for brood year 1966 (Figure 3) when the river fry migrated considerably earlier than those from the channel. Since the river fry were less well developed (i.e. had more yolk) and migrated to a greater extent in the daytime, it is likely that they were being "scoured" from the river gravel by high water velocities. Egg deposition occurred at approximately the same time in both the river and the channel.

The timing of the river fry migration has varied little since 1962 (Table III). The variation which has occurred is probably related to river discharge and water temperatures.




Figure 3. Sockeye fry migration, Fulton River and Channel, 1967.

(b) Fry Quality Studies

It is imperative to determine whether or not the fry produced in spawning channels are of a quality comparable to that of naturally produced fry. The development program will be most successful if the channel fry are able to survive as well as the river fry when placed in the natural environment. A study was carried out in 1967 to compare the quality of river and channel fry using various indices of quality.

(i) Length, weight and development stage

Fifty fry were randomly selected from each population. These were measured to the nearest millimeter, weighed to the nearest milligram and graded for development stage. Development stage gradation consisted of classifying each fish as one of five stages, ranging from alevin to buttoned-up fry, based on the amount of yolk sac visible externally.

The results showed that the river fry were significantly heavier (160 mg. compared to 147 mg.), insignificantly longer (29.4 mm. compared to 29.3 mm.), and less well developed.

(ii) Swimming performance

Samples of 35 fry from each population (selected for development stage) were fin-clipped for later identification and placed


in an experimental flume for a 12 hour conditioning period at water velocities of approximately 0.1 ft/sec. The test procedure consisted of recording the order of dropout (ODO) as the water velocity in the flume was gradually increased.

A percentage difference between the means of the ranked ODO's was calculated (Bams, 1967) and the Mann-Whitney U-test was employed to determine the statistical probability of obtaining such a percent difference.

Pooled results from 57 tests (random and non-random samples) revealed that 47% of the tests statistically favoured river fry (only 14% favoured channel fry).

(iii) Starvation resistance

Non-random samples of each population (i.e. selected for a common development stage, usually buttoned up fry) were placed in separated portions of an aquarium with a filtered recirculating water supply. Periodic subsamples were made to determine the rate of weight loss and the tests were continued until most of the fish had died of starvation (usually about 20 days). In general, the river fish survived significantly longer but appeared to lose weight at a



A view of the interior of the
Department of Fisheries fry quality
laboratory at Fulton River.

slightly faster rate.

(iv) Vulnerability to predation

A number of wild coho salmon smolts and rainbow trout were captured and held in 20 gallon aquaria. These predators were conditioned to their aquaria and periodically fed sockeye fry until they demonstrated an active ability to pursue and capture all available fry within a 12 hour period.

Random and selected samples, varying from 50 to 100 fry per population, were differentially fin-clipped and released into the predator aquaria. The test was terminated when the predator had consumed approximately 50 percent of the prey. The survivors were then removed and identified to source. A chi-square analysis was then carried out to determine whether any differential mortality had occurred. The tests were replicated using single predators and group predators. A control, using an equivalent number of marked fish, provided a check on natural mortality.

A pooled chi-square from 24 tests demonstrated that there was no significant difference between the two fry populations. Only three tests showed any significant

difference in vulnerability to predation and all of these favoured river fry.

(v) Biochemical composition

Fifty fish from each population sample were selected for a common development stage (usually buttoned-up fry). They were measured to nearest millimeter and weighed to the nearest milligram before being sent to the Vancouver Technological Station of the Fisheries Research Board of Canada.

Total nitrogen was determined by the micro-Kjeldahl procedure, total solids by drying the fish to constant weight at 105°C, and total lipid by the method of Folch, Lees and Stanley (1957). Observation of the graphs of log weight on log length, log lipid on log weight, solids on weight, and nitrogen on weight, reveal no differences between channel and river fry. Statistical analysis is continuing.

In conclusion the preliminary results indicate that the river fry were superior to channel fry in terms of weight, stamina and resistance to starvation. The length, biochemical composition and predator avoidance ability of the two fry populations were not shown to be significantly different. It is of interest to note that despite

the apparent inferiority of the channel fry they appeared to survive as well in Babine Lake as did their counterparts from the river (see section III, C, 2).

The fry quality program will be continued in 1968 but some changes in technique are anticipated. Absolute measures of the quality indices (rather than relative as in 1967) will be obtained, so that differences in fry migration timing will not confound the effect of the incubation environment and make statistical analysis difficult.

2. The Distribution, Growth, and Survival of Sockeye Fry Produced from the Fulton River and Artificial Spawning Channel in 1966

This study was designed to answer questions related to the evaluation of the Babine Lake sockeye development program. These are: (1) how do fry produced from a spawning channel compare to those produced from natural spawning grounds in terms of lake distribution, growth and survival; and (2) what part of Babine Lake is used as a nursery area by Fulton River fry?

Sockeye fry were captured during their downstream movement from the Fulton River and the adjoining artificial spawning channel. The fry were marked distinctively by removing the right pelvic fin from the river fish and the left pelvic fin from the channel fish. Both groups were then released to enter

the lake. A total of 203,754 river fry and 299,363 channel fry were marked and released between May 8 and June 23, 1966. Dates of 50% release of marked fish were June 6 for river fry and June 17 for channel fry.

Recoveries of marked fish were made by examining pre-smolt sockeye caught by purse-seining in the main lake area throughout the summer and fall. In addition, marked smolts were recovered at the lake outlet the following spring.

Distribution in the Lake

Figure 4 shows the location of mark recoveries for three periods in 1966, while Figure 5 shows the distribution of the total under-yearling population as indicated by catch-effort data. Fry from Fulton River dispersed rapidly and widely into the main lake area. Most moved southward initially and occupied areas in the southern half of the lake. Later (August-September) a northward shift in distribution was evident and by October young sockeye were most abundant in the lake area adjacent to the Fulton River. Channel fry did not appear to disperse as widely as river fry - probably because lake entry of channel fry was, on the average, later.

Growth

No difference in the relative growth rates of channel and river fish could be detected by




Figure 5. Estimates of the relative density and abundance of marked and unmarked sockeye by lake area and fishing period. See Figure 4 for location of areas.

covariance analysis of mark recovery data obtained throughout the first 4-5 months of lake residence. However, channel fry were smaller, on the average, probably due to late lake entry and thus a reduced growing period.

Survival

Survival rates of channel and river fish were examined by comparing the mark ratio at the time of release as fry (1 river mark to 1.47 channel marks) to the ratios observed in fish caught in the lake in three subsequent sampling periods (June-October). Analysis, using chi-square (Table VI), did not reveal any significant departures from the expected ratio.

Growth and Survival to the Seaward Migrant Stage


During the course of the 1967 smolt run, 487 marked river fish and 790 marked channel fish were recovered from 174,957 smolts examined. Preliminary analysis of the data indicates (1) that channel fish continued to be smaller than river fish (mean lengths of 76.9 and 78.9 mm. respectively) and (2) that differences in survival rate to the smolt stage were probably not statistically significant ($\chi^2 = 2.55, p > 0.10$).

TABLE VI. CHI-SQUARE TESTS OF DEPARTURES FROM EXPECTED RATIO
OF RIVER AND CHANNEL RECOVERIES, 1966

Fishing Period	Marks Recovered			Ratios		Chi-Square
	River	Channel	Total	Expected	Observed	
I. June 25-July 27	35	62	97	1:1.47	1:1.77	0.69
II. August 16-September 9	52	75	127	1:1.47	1:1.44	0.03
III. October 6-25	23	28	51	1:1.47	1:1.22	0.32

1
32
1





Employees of the Fisheries Research Board
of Canada clipping pelvic fins of sockeye
fry in the laboratory at Fulton River.

D. Primary Productivity of Babine Lake

The rationale behind the Babine development program is that the main lake basin, on the average, rears only one-fifth as many sockeye as the North Arm - Nilkitkwa area while it is potentially as productive. In 1966, relative primary productivity of the various basins of Babine Lake was assessed utilizing the C^{14} in situ method. It was found that primary productivity was slightly lower in the North Arm (outlet) and in Morrison Arm (an inlet) than in the main lake region (Narver, 1967). An unusually high rate of photosynthesis at one part of the main lake in September may have been related to the decomposition of salmon carcasses in a nearby stream.

Morrison Lake, a major tributary, was much lower in rate of photosynthesis, pH, alkalinity, compensation depth, and total dissolved solids than any part of Babine Lake. The rate of carbon fixation of Babine Lake was much less on a per unit area basis than those reported by other workers for the majority of 24 sockeye salmon lakes similarly tested in Southwestern Alaska but was similar to most per unit volume of the euphotic layer. Much of this difference is attributable to a shallower euphotic zone in Babine Lake. Ranked with the 24 Alaskan lakes, Babine was first in alkalinity, third in TDS, and fifth in pH.

The productivity measurements suggest that the carrying capacity for sockeye salmon per unit of nursery area of the main lake region is at least as great as that of the North Arm.

E. Other Studies

1. Adult and Smolt Sockeye Enumerations

Among the measurements needed for assessment of the Babine Lake development project are adult escapements to Babine Lake and the smolt runs resulting from them. The relationship between these two parameters will serve as one measure of the success of year classes and may be quite sensitive to changes in the balance between the rearing capacity of the lake basins and the numbers of young introduced into them. Ultimately, a measure of spawning escapement and resulting smolt output from particular production units or stocks is needed and some progress is being made in this regard. Since 1946 total adult escapement has been counted annually at the Babine fence and spawning populations have been counted or estimated in individual streams. Tagging programs in 1958, 1962 and 1963 have demonstrated the approximate times of passage of adults of these particular stocks at the fence.

Populations of sockeye smolts migrating out of Babine Lake have been assessed since 1951: the "late run" produced by spawners in the main lake basins, 1951-1957; the entire run - including North Arm - Nilkitkwa production - after 1957. Assessment of reproductive success of individual stocks, however, will require better information than now exists on the

timing of their smolt migrations. At the present time it is only possible to assess production from two groupings of fish, those contributing to the early and to the late runs.

Fin-clipped smolts of the Fulton River stock were recovered from the seaward migrations of 1967 and in 1968 and 1969 smolts from subsequent markings will be available. It is expected that the timing of this stock will become better known. Similar information for other stocks may be gained in the same way.

Counts of adult Babine sockeye passing through the fence and in the various streams for the years 1946-1966 are shown in Table VII. The relationship between potential egg deposition (calculated from Table VII and fecundity counts) and smolts produced is shown in Figure 6 for three groupings:

1. the entire system - assuming all females entering the lake spawned;
2. the North Arm - Nilkitkwa area; and
3. the main lake area including Morrison Arm.

The main lake and North Arm regions are calculated from a stream count base because differences in fence counts and summed stream counts cannot be safely attributed to a specific region. In each region there is a general trend but again a large variability in production per million eggs potential is apparent.

TABLE VII. BABINE SOCKEYE ESCAPEMENTS, IN THOUSANDS OF FISH, 1949-1966.

	1949	1950	1951	1952	1953	1954	1955	1956	1957	1958	1959	1960	1961	1962	1963	1964	1965	1966	1949-1966	
																			Sums	Means
Babine fence count	461	364	141	349	687	494	71	355	433	812	783	263	942	548	588	828 ¹	580	389	9088	504.9
Indian catch	29	27	19	34	27	22	10	31	20	39	17	17	32	18	20	20	19	19	420	23.3
Potential spawning stock (count less catch)	432	337	122	315	660	472	61	324	413	773	766	246	910	530	568	808	561	370	8668	481.6
<u>North Arm - Nilkitkwa Region</u>																				
Upper Babine River	216.0	65.0	13.3	78.2	147.0	136.7	9.7	66.5	117.8	156.8	156.7	36.9	196.0	192.0	119.3	222.0	120.4	69.0	2119.3	117.7
Lower Babine River	135.0	116.0	10.8	69.0	127.4	100.0	9.0	52.3	66.5	107.8	123.5	54.0	171.5	61.0	34.5	46.0	176.0	114.0	1574.3	87.5
9 Mile	0.9	1.0	0.4	0.1	2.5	1.0	0.1	0	4.0	0	2.4	1.8	2.5	0.5	1.0	1.5	0.5	0.8	21.0	1.2
5 Mile	0	0.1	0.1	0	0.3	0.3	0.1	0	0.2	0	0.6	0	0.5	0.1	0	0.1	0.2	0.2	2.8	0.2
Total spawners	351.9	182.1	24.6	147.3	277.2	238.0	18.9	118.8	188.5	264.6	283.2	92.7	370.5	253.6	154.8	269.6	297.1	184.0	3717.4	206.5
Per cent of potential spawners	81.5	54.0	20.2	46.8	42.0	50.4	31.0	36.7	45.6	34.2	37.0	37.7	40.7	47.8	27.3	33.4	53.0	49.7		42.9
<u>Main Lake Region</u>																				
Morrison system	1.6	5.9	4.1	1.2	24.7	24.0	1.8	27.0	28.9	18.0	35.9	9.9	23.6	12.5	41.8	27.0	8.5	8.8	305.2	17.0
Fulton River	33.9	42.0	15.2	31.5	134.4	105.6	16.7	81.0	108.0	76.0	114.0	36.0	170.1	86.4	98.6	117.0	123.3	59.1	1448.8	80.5
Pinkut Creek	10.5	12.0	4.9	7.5	23.5	25.0	3.2	22.8	29.1	44.0	77.6	27.0	44.1	21.4	40.0	135.3	22.0	21.4	571.3	31.7
Pierre Creek	4.2	17.9	11.5	3.3	19.2	17.0	3.2	18.0	21.2	29.4	33.0	9.9	24.5	4.1	28.4	22.0	10.0	8.8	285.6	15.9
Grizzly	1.5	2.7	2.1	3.5	6.0	3.1	0.5	4.8	7.0	30.0	14.0	10.8	23.5	4.6	11.4	8.0	5.0	4.5	143.0	7.9
Twin	2.3	7.6	4.8	0.4	9.8	14.0	2.4	4.5	5.4	12.0	9.0	5.4	6.9	1.3	11.4	9.0	3.0	2.0	111.2	6.2
4 Mile	1.6	4.2	0.9	0.2	2.0	2.2	0.4	0.4	2.5	6.0	5.4	1.8	1.0	2.8	2.8	2.5	1.4	1.7	39.8	2.2
Tachek	2.6	2.6	2.5	0	2.4	1.9	0.3	0	6.4	1.8	6.0	1.8	0	0.6	1.6	3.0	0.7	0.3	34.5	1.9
Sockeye	0.2	0.9	0.8	0	0.6	0.9	0.5	0	2.5	1.5	4.0	1.8	0	1.0	2.4	1.5	0.1	1.4	20.1	1.1
6 Mile	0.4	1.2	0	0	2.6	1.8	0.1	0.1	0.6	2.3	3.5	0.9	0	0.9	1.4	1.5	0.1	0.3	17.7	1.0
Pendleton	1.1	1.2	0	0	1.4	1.1	0	0	0.3	0	2.5	0	0	0.2	0	1.4	0	0	9.2	0.5
Others ²	0	0	20.0	74.4	1.0	0	0	0	0.2	72.5	3.9	0.3	51.8	6.2	6.2	9.3	1.8	0	247.6	13.7
Total spawners	59.9	98.2	66.8	122.0	227.6	196.6	29.1	158.6	212.1	293.5	308.8	105.6	345.5	142.0	246.0	337.5	175.9	108.3	3234.0	179.7
Per cent of potential spawners	13.9	29.1	54.8	38.7	34.5	41.7	47.7	49.0	51.4	38.0	40.3	42.9	38.0	26.8	43.3	41.8	31.4	29.3		37.3
Total spawners accounted for	411.8	280.3	91.4	269.3	504.8	434.6	48.0	277.4	400.6	558.1	592.0	198.3	716.0	395.6	400.8	607.1	473.0	292.3	6951.4	386.2
Per cent accounted for	95.3	83.2	74.9	85.5	76.5	92.1	78.7	85.6	97.0	72.2	77.3	80.6	78.7	74.6	70.6	75.1	84.3	79.0		80.2
Potential spawners not accounted for	20.2	56.7	30.6	45.7	155.2	37.4	13.0	46.6	12.4	214.9	174.0	47.7	194.0	134.4	167.2	200.9	88.0	77.7	1716.6	95.4
Per cent not accounted for	4.7	16.8	25.1	14.5	23.5	7.9	21.3	14.4	3.0	27.8	22.7	19.4	21.3	25.4	29.4	24.9	15.7	21.0		19.8

¹Estimate derived from stream counts, tag and recovery, av. "not accounted for" 1949-1963.

²Includes: a intermittent counts in small marginal streams

b counts of fish which died unspawned esp. 1951, 1952, 1958, 1961

c for Nanika egg take from Pinkut Creek; 1961 = 2050, 1962 = 6200, 1963 = 6200, 1964 = 9300, 1965 = 1800




Figure 6. Babine Lake sockeye smolt production.

The 10 years of production data for the entire system suggests that some increase in smolts has usually accompanied increases in potential egg deposition, but they also reflect a great deal of variability in production at all levels. Brood years 1958 and 1961 had similar egg depositions but there was a fourfold difference in smolt production; brood years 1960, 1961 and 1965 had potentials ranging from 400 to 1600 million eggs but produced similar-sized smolt runs.

Factors Governing Age at Maturity

Management of sockeye stocks is hampered by the difficulty of forecasting runs well in advance of commercial fishing seasons. Part of this problem is caused by the diverse and variable age at maturity of the species.

Two lines of investigation are being followed at Babine. First, sampling of the individual production units is providing a basis for study of parent-progeny age relationships. Figure 7 shows the age composition of these units for the years 1962-1966. Second, studies at Four Mile Creek on Babine Lake have shown that mating is not random with respect to the age composition of the fish present (Hanson and Smith, 1967). There are two factors which may limit the participation of 4-year-old males in the reproductive act. There is a strong tendency for sockeye of like size to mate,

AGE COMPOSITION — BABINE SOCKEYE

(MALES and FEMALES COMBINED)

1-2 AGE SPAWNER — 1-3 AGE SPAWNER • N = LESS THAN 30




Figure 7. Age composition - Babine sockeye.

i.e. 5-year-old males mate with 5-year-old females, and 4-year-old males mate with 4-year-old females. On occasion the 4-year-old males are displaced by the larger 5-year-old males and relegated to a "satellite" status, but the inverse is seldom true. These factors limit contribution of 4-year-old males to the population gene pool and could have a marked effect upon the ultimate ages of progeny if heritable factors are involved.

The age composition of sockeye stocks in other Skeena tributary systems (Table VIII) has been assessed whenever possible as an additional step in the search for cause and effect relationships in age at maturity studies.

Smolt Size and Post-Lacustrine Survival

Efforts to obtain efficient utilization of lake nursery areas and the forecasting of adult returning population will be aided by an understanding of the relation between smolt size and ocean survival. Both the numbers and the age composition of returning sockeye may be adversely affected by increasing the young salmon population beyond some optimum level.¹

¹ It has been estimated that the lake nursery area could support about 350 million more fry than can be produced from existing spawning grounds, but it will be necessary to learn what numbers of fry will produce smolts of a size and quality likely to provide the largest fishable return.

In 1966 and 1967 smolts in five separate size categories were marked by Bergman-Jefferts coded wire tags. Recoveries in the commercial fishery, at the Babine fence and on the spawning grounds will be used to measure marine survival rates. In 1966, 47,000 smolts were tagged and four tagged jacks were recovered in the escapement of 1967.

Survival of certain sizes of smolts may at times be influenced by parasitism. The tapeworm Eubothrium salvelini is associated primarily with smolts of small size (Dombroski, 1955; Margolis and Boyce, 1963) and was present in all daily samples in 1966 and 1967 (Figure 8). Parasitized smolts succumb to routine handling more readily than unparasitized smolts and their scale patterns differ. The relation between this parasite and survival of sockeye is not yet known but is under study.

2. Factors Associated with Age at Maturity of Skeena River Sockeye

Several correlated effects have led to a possible chain of events which may influence the age at maturity of Skeena River sockeye salmon. These are:

1. in general the size of 1.3 females is larger than that of 1.2 females;
2. a significant positive correlation exists between body size and egg size, hence eggs of 1.3 females are, on the average, larger than those of 1.2 females;




Figure 8. Parasitism of Babine sockeye smolts by tapeworm Eubothrium salvelini in 1966 and 1967.

3. results from reciprocal crosses of 1.2 and 1.3 sockeye indicate that size of resulting alevins and their subsequent growth as juveniles is positively correlated with egg size, and that this is a maternal effect;
4. within each of four brood years of Skeena River sockeye, fish maturing as 1.2's had a higher number of freshwater circuli than those which matured as 1.3's;
5. number of freshwater circuli is directly correlated with smolt size.

The evidence thus suggests that the size of the female parent may directly influence the age at which the progeny will mature. This further suggests that, where other influencing factors are equal, an alteration of generation time can exist, e.g. that 1.3 females will produce 1.2 females which will in turn produce 1.3 females. This hypothesis is now being tested.

3. A Review of the Literature on the Effects of Changes in Temperature Regime on Developing Sockeye Salmon Eggs and Alevins

(a) Introduction

At a meeting of the Babine Lake Sockeye Development Sub-Committee, February 21, 1967, concern over a possible 10-15^oF rise in temperature under certain conditions between the inlet and outlet points of the Fulton spawning channel was

expressed. The biological consequences of temperature alteration in artificial spawning channels were not known to this group and a survey of literature was to be undertaken.

This report specifically deals with the effects of alteration or variation of temperature from optimal conditions for larval development of sockeye salmon. It is beyond the scope of this review to speculate about changes in temperature regime which might be expected in a particular facility. Little of the available literature is specifically directed to the species in question hence it is necessary to apply generalities resulting from studies on a broad spectrum of species. Excellent reviews regarding the effects of temperature on development are already available: Hayes (1949), Seymour (1956), Rockwell (1956) and Kinne and Kinne (1962). A large segment of the applicable literature has been reviewed for this report, but the survey was not exhaustive. It is believed, however, that the more essential aspects are covered, and that little would be gained from further searching. Because many physiological processes, such as rate of growth, are influenced by more than one environmental factor, many of which interact (e.g. O_2 and temperature), treatment of the effects of temperature

alone is not entirely satisfactory. For the immediate purposes of this report this simplification appears necessary.

There are many aspects to a particular temperature regime, such as the mean value, maxima and minima, the order and frequency of fluctuations, the rate of change, the duration at a certain level, etc., which can directly affect an organism. Not only may any one aspect influence one or more biological processes or events, but biological effects may be accumulative or interdependent. In general, effects of a temperature change are many, and extremely varied. They can range from a simple, direct relationship, such as the accelerating influence of an increase of temperature (within a certain range) on the basic metabolic rate, to extremely complex and multi-order effects. For example, the final effect of a rise in temperature on the survival potential of a particular migrant fry may go through the following chain of events: (a) an increase in the numbers of aerobic organisms in part of the channel; (b) a subsequent reduction in O_2 -concentration; (c) mortality of part of the brood; (d) decomposition of these dead larvae by microorganisms; and (e) pollution of the environment in which this fry migrant developed. The final outcome may be death or lowered viability.

Clearly effects of this nature will be determined by local conditions and cannot be predicted in advance. However, they are of importance as they affect the environment and hence may limit the population.

(b) Effects During Period Before Egg Deposition

Pacific salmon spawn during the declining phase of the annual temperature cycle. At Babine, sockeye generally spawn at temperatures between 55° and 45°F. If the decline in temperature, after summer maximum, is delayed a similar delay may be expected in spawning (Sheridan, 1962). Some immediate effects may be: (1) initiation of development of the embryos is delayed; (2) the sexual products become "overripe" and reduced egg survival results, particularly through loss of eggs (before fertilization) and reduced fertilization due to reduced viability of either eggs or sperm, or both. Extensive prespawning mortality in sockeye adults has been correlated with high water temperature (Royal, 1953). From Brett's work (1952, 1967) on thermal resistance in sockeye salmon, the zone of thermal resistance is known; generally speaking, temperatures become lethal at around 25°C (= 77°F). However, temperatures between the optimum level at around 15°C (= 59°F) and the upper lethal level become increasingly

stressing and the scope for further activity is reduced, as evidenced by a gradual decrease in swimming performance (Brett, 1967). Further consequences of temperature levels above the physiological optimum can be expected, e.g. a reduced life span due to accelerated metabolism, and higher incidence of disease, particularly attack by Saprolegnia spp.

(c) Effects During the Larval Stages

Changes in sensitivity during development

Not all larval stages are equally sensitive to similar amounts of stress. In relation to O_2 and temperature the most sensitive periods are the period between early cleavages to the closing of the blastopore, and the period immediately prior to hatching (cf. Battle, 1944; Hayes and Armstrong, 1942; Hayes, 1949; Garside, 1959; Swift, 1965). Apparently both heat and cold tolerance of developing fish larvae is limited at the early stages and gradually increases during development (e.g. Donaldson, 1955, and Olson and Foster, 1956).

Rates of differentiation and growth (development) and metabolism

Developing sockeye salmon larvae depend entirely on their yolk supply as a source of

energy during the full period of larval development. This supply must satisfy all physiological needs, i.e. various additive demands must share the available reserve. During normal development a rather precise balance is struck between the competing demands for activity, metabolism, and growth. A fully developed fry is formed at a time, stage of development, and size, optimally adapted to the average conditions experienced by the stock during succeeding stages. If, due to unusual circumstances during development, the demand of any one of the competing functions increases, one or both of the others must suffer if the supply rate remains constant, or, alternatively, the developing larva will use up its supply before development is complete.

The effect of temperature alone on the larval rate of development is an important consideration. In general a rise in temperature increases the rate of development, thus decreasing the developmental time period. Developmental rate is expressed as the reciprocal of developmental time. The relationship between this rate and a variety of constant temperatures is nearly linear over most of the normal developmental range, and

such an interpretation forms the basis for the commonly used constant "Day-Degree" rule used by most investigators (Hayes, 1949; Seymour, 1956; Rockwell, 1956). These authors have extrapolated the linear approximation to obtain an estimate of the temperature at which developmental rate is zero. Over extended temperature ranges, however, the relationship is definitely curvilinear, and such extrapolations are in error.

Experimental evidence for developmental rates at temperatures below 2.5°C is scarce. Embury's (1934) work on Salvelinus fontinalis and Salmo trutta, however, clearly demonstrates a gradual slowing in deceleration of the growth rate with decreasing temperature at these low levels.

Garside's data for lake trout (Garside, 1959) have been selected to illustrate the changes in rate of development at lower temperatures. Garside ignored the non-linearity of his data, and indicated a development rate (100/(days from fertilization to hatching)) at 0 C of between 0.2 and 0.3, corresponding to a time period of some 380 days. From his data a decelerating rate of 13 days per 2.5 C decrease is estimated and,

extrapolating, the rate of development would be 0.503 at 0 C, corresponding to a period of 198 days. Table IX expresses expected increases in rate of development, starting with 0 C for increments of 2.5 C.

With this table the advance in days caused by a given temperature rise over a given number of days can be accurately determined. For example: with the river at 0 C and the channel at 2.5 C during a 20-day period, development in the channel is advanced by 41%, equivalent to 8.2 days. These 8.2 days, however, are based on a developmental rate of 0.503. In the spring when emergence occurs, the temperature is about 5 C, and at this temperature the developmental rate is about 1.031. Thus the early gain of approximately 8.2 days can be made up in about four days ($8.2(503/1031)$) by the river fry at the higher temperature. Emergence of the two groups of fry may occur at the same time because of factors other than temperature, e.g. daylight, length and silt, however the relative stages of development of the two groups will have been altered. The developmental rates for many salmon species are quite similar and the lake trout data

TABLE IX. EXPECTED RELATIONSHIP BETWEEN DEVELOPMENT RATE,
FERTILIZATION TO HATCHING, AND TEMPERATURE

Temperature (°C)	Days	Rate (100/da)	Rate Change Over Temperature Interval - Δ Rate and Percent Increase -							
			From 0.0 to		2.5 to		5.0 to		7.5 to	
			Δ	%	Δ	%	Δ	%	Δ	%
0.0	198	.503	-	-	-	-	-	-	-	-
2.5	141	.709	.206	41.0	-	-	-	-	-	-
5.0	97	1.031	.528	105	.322	45.5	-	-	-	-
7.5	67	1.493	.990	197	.784	111	.462	44.8	-	-
10.0	50	2.000	1.497	298	1.291	182	.969	94.0	.507	34.0

Note: The maximum relative increase in rate occurs around 5C.

compare favourably with those available for sockeye at higher temperatures (Rucker, 1937).

Table X gives the estimated development rates between 0 and 10 C in 0.5° intervals. From it rate changes can be calculated for any temperature interval according to the method shown in Table IX.

TABLE X. DEVELOPMENT RATES AT DIFFERENT TEMPERATURES

Temperature °C	Rate (100/days)	Temperature °C	Rate (100/days)
0.0	.503	5.5	1.111
0.5	.534	6.0	1.198
1.0	.570	6.5	1.293
1.5	.611	7.0	1.392
2.0	.657	7.5	1.493
2.5	.709	8.0	1.594
3.0	.765	8.5	1.695
3.5	.825	9.0	1.796
4.0	.889	9.5	1.898
4.5	.958	10.0	2.000
5.0	1.031		

Note: For rate values located between values listed use linear interpolation.

Admittedly these calculations are somewhat more involved than the Degree-Day method, but the greatly increased accuracy,

particularly at the low end of the temperature range, dictates its use.

From the rates shown it is evident that even limited changes in temperature regime have a significant effect on the developmental rate. Hence, a consistently higher daily average temperature will result in earlier completion of development. Such a condition may lead to two alternatives: (1) early emergence and migration at a time that will not provide optimum conditions for further growth and development; or (2) remaining in the gravel with further depletion of critically low energy reserves, leading to poor condition at time of emergence. Both alternatives are considered undesirable: "premature emergence in good condition" because the lake environment (e.g. temperature and/or food) is not yet favourable, and "suboptimal condition at the right time" because such fry have been shown to exhibit a reduced survival potential (Bams, 1967).

Other effects

In addition to alteration of the developmental period, changes in temperature affect differentiation and growth in other aspects. A decrease in potential size of fry

at the end of the incubation period from increased temperature is well documented (cf. Gray, 1928; Donaldson, 1955; Seymour, 1956). Gray attributes the reduction in size to differences in rates of reaction as these pertain to growth and maintenance: relatively more energy is required for the increased rate of maintenance metabolism, hence less yolk is available for growth. Sockeye fry of a reduced size are known to have a decreased viability as shown by Brett (1952) for low temperature tolerance, and by Bams (1967) for swimming performance and vulnerability to predation. It is now thought likely that such changes will lead to an alteration of life history and time of maturity.

Related effects are changes in morphological characteristics of the developing larvae. There are a number of critical stages during development when different levels of O_2 and/or temperature can affect the extent of various differentiation processes. Particularly the work of Hayes (cf. Hayes, 1949; Hayes et al., 1953) has shown that various differentiation processes have different temperature coefficients so that interacting processes affect each other differently at different temperatures.

Much evidence is on hand regarding the effect of temperature level on the number of vertebrae, fin rays, morphometric relationships, etc. (cf. Alderdice et al., 1958; Garside, 1959; Kinne, 1963). Seymour (1956) indicates, for chinook fry, changes in vertebrae counts, dorsal fin rays, and anal fin rays. Related to these effects are the influence of temperature on the number of abnormal (deformed) fry and on hatching efficiency. From the limited body of information reviewed it can be concluded that changes will occur with changes in temperature level. Optimum values are indicated and temperature changes precipitate modifications of these characteristics in either direction.

Effects of substantial decreases in temperature are presumed to be limited to events that can occur during a matter of hours only. Direct lethal effects are perhaps unlikely. Brett (1952) and Brett and Alderdice (1958) investigated the lower lethal temperature limits of young chum and sockeye salmon. Their results indicate that exposure to 0 C is within the tolerance range of both species from the fry stage on. Exposure to temperatures far enough below zero to be

harmful (<-0.5 C) cannot occur in fresh water unless the intergravel water freezes. Survival under the latter conditions, even for limited exposure, is unlikely. Direct effects on morphological characters may possibly occur in cases where the duration of the critical phase of a differentiation process is sufficiently short to be influenced by a temporary cold wave. No data were found regarding the duration of any such critical phases.

Continuous low temperature, especially during early embryonic stages, is known to be harmful and to increase mortality. Brett and Alderdice, 1958, mention that fry and yearlings acclimated to 2.5 C showed less resistance to cold (below 0 C in salt water) than did fry acclimated to 5.0 C, and infer that the 2.5 C acclimation temperature imposed a stress on the fish.

The dependence of the effect of a change in temperature on previous thermal experience is well known from temperature tolerance work by various authors. In general, exposure to high or low temperature can be tolerated better if the fish is already acclimated to a high or low temperature level. The effect is much

more pronounced (greater dependency) for the low than for the high temperature limit. Hence, a sudden decrease in temperature is potentially much more harmful than a slower decrease of the same magnitude; the rate of change of an increase in temperature is of much less importance.

The effects of temperature fluctuations are largely unexplored. As long as harmful extremes are not reached little effect of increased fluctuation around a common mean can be expected. In regard to morphological characteristics Lindsey and Ali (1965) established, in a temperature transfer experiment with medaka, that these characteristics followed the arithmetic mean temperature. It should be noted that, due to the previously mentioned reduced tolerance ranges during certain stages in larval development, increased daily fluctuations of a certain magnitude could be harmless during most of the developmental period but potentially harmful at certain times, particularly during the pre-eyed stages.

Indirect effects

An immediate effect of increased metabolic rate due to temperature increase is

increased O_2 demand. The importance of O_2 to the brood and effects of changes in demand and/or supply are discussed below.

Most indirect effects of a change in temperature are expected to occur by means of changes in the environment. Both changes in physico-chemical and biotic elements of the environment are likely to occur. Of the former oxygen is of major concern. The direct influence of available O_2 on the fry is of prime importance, possibly more so than temperature per se. O_2 is known to affect differentiation rate, growth rate, efficiency of yolk conversion, mortality rate at all embryonic stages, hatching success, time of emergence, formation of abnormalities and morphological characteristics. Temperature affects the O_2 content of the water directly by reducing the saturation level as temperatures increase. Indirectly O_2 content will be affected through increased consumption by the resident flora and fauna. Conversely, a lowering of the temperature could cause over-saturation, which is very harmful to fish, and increased ice formation, which could seriously interfere with the intergravel water flow pattern. Of the biotic factors possible,

increases in algal and bacterial growth as correlates of increased temperature should be mentioned, both of which would have consequences regarding O_2 content, metabolic waste products, and possibly interference with the intergravel flow. Incidence and growth of fungus on dead eggs can also be expected to increase, with a subsequent higher mortality rate of adjacent eggs.

The effects of lowered O_2 content and reduced flow have been studied extensively by Garside (1959, 1966), Silver et al. (1963), and Shumway et al. (1964). Small reductions in either temperature or O_2 were shown to measurably affect the fry, especially in respect to rate of development and size, and increasingly so as the absolute levels of these factors dropped. Of considerable interest here is the repeatedly established interaction of O_2 and temperature (Alderdice et al., 1958; Garside, 1959, 1966). A progressive increase of any effect (such as retardation of development rate) produced by a certain level of hypoxia was observed with increase of temperature. Also, with increasing temperature the effect occurs at progressively earlier stages in development, i.e. a certain level of

hypoxia may not adversely affect the developing larva of a particular stage at a certain temperature, but it will do so at: (1) a later stage at the same temperature, and (2) at its present stage if the temperature is increased. These findings illustrate the earlier observation that the consideration of the effects of a single factor are not satisfactory, and that an integrated multi-factor model is necessary to assess realistically the effects of any one factor, such as temperature.

Conclusions

1. The effects of changes in temperature regime on developing salmonid larvae are of great diversity, and many are potentially harmful.
2. Virtually all the effects mentioned are impossible to quantify beforehand because of (a) the body of data applicable to the stock under consideration is too limited, and (b) the extent of the expected temperature changes is unknown. Prediction of changes in second, or higher order events that are likely to affect the brood is also impossible due to the many,

largely unknown, local conditions that will influence these events.


4. Diel Vertical Movement of Pelagial Sockeye Salmon Juveniles

A better understanding of the sockeye's activities is needed to determine their requirements for growth and survival and to determine the capacity of different parts of the lake to meet these requirements. In 1967, attention was directed toward the diel movements of the sockeye and their food supply.

(a) Problem and Methods

A general pattern of vertical diel movement of pelagial sockeye salmon juveniles during the summer in Babine Lake was suggested from tow netting (1956 to 1963) and from echo sounding, tow netting and purse seining in 1966. These sources indicated that (1) during the day juveniles were deeper than 50 ft., (2) at least some moved to the lake surface at dusk, (3) at least some remained within 50 ft. of the surface at night, and (4) juveniles moved away from the surface soon after dawn. The objective of this research in 1967 was to answer specific questions about this apparent diel vertical movement of juveniles: "What is the precise pattern of movement? When does this behaviour first become evident and how does it change through the summer? Is this behaviour the

CANADA
FISH
BO
LINE



The Fisheries Research Board of Canada
purse-seining for sockeye salmon in Babine Lake.

same in the North Arm as in the main lake? What is the physical and biological environment associated with the behaviour? What is the feeding biology of juveniles and how does it fit into this behaviour?" The basic hypothesis was that juveniles seek a specific light intensity and hence light is the primary factor controlling the diel movement. To test this a prototype underwater light meter was designed and ordered, but unfortunately was not delivered in time for field use in 1967.

The field study was based on observations and samples taken at intervals during ten 24-hour series (noon to noon). These series included transects with a 145 kc Sea Scanar (Honeywell) echo sounder, tows with a 3-ft. Isaacs-Kidd midwater trawl, purse seine sets with a 600 x 35 ft. net, horizontal zooplankton tows with Miller samplers at eight depths, secchi depth, and vertical temperature profile. These 24-hour series were conducted from mid-July to early October in the North Arm of Babine Lake (7) and in the main Babine Lake (3).

(b) Early Results

The success of this investigation lay primarily with the reliability of the echo sounder as a device to indicate the abundance of juvenile sockeye salmon at any depth. A correlation coefficient of .81 was obtained in the relationship

between juvenile catch by trawling and the number of echo sounder targets (Figure 9). Purse seine sets substantiated this since midday sets when there were no echo sounder targets above 35 ft. resulted in no catches of juveniles; sets made in the evening, night, or early morning, when targets were found above 35 ft., resulted in catches of juveniles.

A precise, well-defined diel vertical movement is displayed by pelagial sockeye juveniles in Babine Lake from at least mid-July to September (Figure 10). The maximum vertical movement in the North Arm was from the midday depth on the bottom or 130 ft. to the lake surface just after sunset. (In the study area between Nine Mile and Five Mile Creeks depth of bottom ranged from 100 to 170 ft.) A distinct and consistent double layer of targets, believed to be juveniles, was found during the day in the North Arm (Figure 10). The two layers maintained a constant distance apart until the top layer reached the surface in the evening. The bottom layer (the majority of targets) continued to rise until it also reached the lake surface (Figure 10). The pattern of diel vertical movement changed in the fall as indicated by the October 2-3 series. Juveniles no longer came to the surface but remained in midwater through the night and




Figure 9. Plot of midwater trawl catch per minute of juvenile sockeye and the simultaneous number of echo sounder targets per minute in the 10-ft. stratum of water at the depth of the trawl. Data from August, 1967 in the North Arm of Babine Lake.




Figure 10. Echo sounder targets per minute (.3 mile) by 10-ft. strata during the 24-hour series of August 5-6 in the North Arm. (Note that time is not to scale.)

returned to deeper water during midday (Figure 11). The double layer was no longer conspicuous during the day.

The July and August diel vertical behaviour of juveniles in the main lake was similar to that of the North Arm except the midday maximum depth was 150 to 170 ft.

The zooplankton of Babine Lake is composed of the Calanoid copepods Diaptomous (2 species), Epischura and Heterocope; the Cyclopoid copepod Cyclops; and the Cladocera Daphnia, Bosmina, and Holopedium. Only Bosmina and Heterocope displayed a pronounced diel movement and these were in opposite directions (Figure 12). Daphnia, Holopedium, Epischura, and both Diaptomous were found almost entirely above 35 feet. Cyclops occurred in greatest concentrations below 35 feet (Figure 12). The species are ranked in relative abundance from lowest to highest: Holopedium, Epischura, Heterocope, Bosmina, Daphnia, Diaptomous pribilofensis, Cyclops and Diaptomous ashlandi. Although Heterocope was relatively scarce it formed a major portion of the zooplankton biomass since an individual adult is about 4 mm. long.

A preliminary examination has been made of juvenile stomach contents from one 24 hour series in North Arm. These data suggest that intensive




Figure 11. Echo sounder targets per minute (.3 mile) by 10-ft. strata during the 24-hour series of October 2-3 in the North Arm. (Note that time is not to scale.)




Figure 12. The diel vertical distribution of zooplankton species in the North Arm of Babine Lake on September 22-23, 1967. Each depth is expressed as a percent of the number of that species taken in a particular vertical profile.

feeding occurred at dusk as the juveniles reached the 0-35 foot stratum and again at dawn as the fish commenced to move away from the lake surface.

There appears to have been no feeding during the hours of darkness when the fish were within 50 feet of the surface and little feeding in midday when the fish were mainly below 100 feet. The dominant food item was Heterocope followed by Daphnia.

IV. ACKNOWLEDGEMENTS

The biological data presented in this report for Fulton River and Pinkut Creek is the result of field surveys conducted by a number of Department of Fisheries personnel, including biologists C. E. Walker, G. A. C. Wilson, D. E. Marshall, R. N. Palmer, M. Giles, R. D. Humphreys, R. K. Kearns, and biological technicians G. A. Buxton, I. A. MacLean and A. P. Stefanson.

Mr. R. K. Kearns submitted the section on the fry quality studies at Fulton River.

The engineering construction section was contributed by Mr. B. A. Heskin. Mr. A. F. Lill contributed a section on the hydrometeorological program.

Dr. W. E. Vanstone was in charge of the biochemical analysis, carried out by Miss A. Oishi.

Scientists of the Fisheries Research Board of Canada submitted summaries of their studies. R. A. Bams contributed the literature review on the effects of temperature on sockeye eggs and alevins, J. McDonald the section on the distribution, growth, and survival of sockeye fry in Babine Lake. H. D. Smith contributed the section on adult and smolt sockeye enumerations, and T. Bilton the section on factors associated with age at maturity of Skeena River sockeye. Dr. D. W. Narver contributed the sections dealing with the primary productivity of Babine Lake and the diel vertical migration of sockeye juveniles.

The Progress Report was edited by L. M. Dill.

V. LITERATURE CITED

- Alderdice, D.F., W.P. Wickett and J.R. Brett, 1958. Some effects of temporary exposure to low dissolved oxygen levels on Pacific salmon eggs. J. Fish. Res. Bd. Canada, 15:229-250.
- Bams, R.A., 1967. Differences in performance of naturally and artificially propagated sockeye salmon migrant fry, as measured with swimming and predation tests. J. Fish. Res. Bd. Canada, 24:1117-1153.
- Battle, H.I., 1944. The embryology of the Atlantic salmon. Can. J. Res., D, 22:105-125.
- Brett, J.R., 1952. Temperature tolerance in young Pacific salmon, genus Oncorhynchus. J. Fish. Res. Bd. Canada, 9:265-323.
- Brett, J.R., 1967. Swimming performance of sockeye salmon in relation to fatigue time and temperature. J. Fish. Res. Bd. Canada, 24:1731-1741.
- Brett, J.R., and D.F. Alderdice, 1958. The resistance of cultured young chum and sockeye salmon to temperatures below 0°C. J. Fish. Res. Bd. Canada, 15:805-813.
- Combs, B.D., 1965. Effect of temperature on the development of salmon eggs. Prog. Fish-Cult. 27:134-137.
- Combs, B.D., and R.E. Burrows, 1957. Threshold temperatures for the normal development of chinook salmon eggs. Prog. Fish-Cult. 19:3-6.
- Dombroski, E., 1955. Cestode and nematode infection of sockeye smolts from Babine Lake, British Columbia. J. Fish. Res. Bd. Canada, 12(1):93-96.
- Donaldson, J.R., 1955. Experimental studies on the survival of the early stages of chinook salmon after varying exposures to upper lethal temperatures. M.Sc. thesis, Univ. Wash. 116 pp.
- Embody, G.C., 1934. Relation of temperature to the incubation periods of eggs of four species of trout. Trans. Am. Fish. Soc. 64:281-292.
- Fish, F.F., and R.E. Burrows, 1939. Notes on the effect of low temperature upon eyed eggs. Prog. Fish-Cult. 45:28-31.

- Folch, J., M. Lee, and G.H.S. Stanley, 1957. A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* 226:497-509.
- Garside, E.T., 1959. Some effects of oxygen in relation to temperature on the development of lake trout embryos. *Can. J. Zool.* 37:689-698.
- Garside, E.T., 1966. Effects of oxygen in relation to temperature on the development of brook trout and rainbow trout. *J. Fish. Res. Bd. Canada*, 23:1121-1134.
- Gray, J., 1928. The growth of fish. II. The growth-rate of embryo of Salmo fario. *J. Exp. Zool.* 6:110-124.
- Gray, J., 1928. The growth of fish. III. The effect of temperature on the development of the eggs of Salmo fario. *J. Exp. Zool.* 6:125-130.
- Hanson, A.J., and Howard D. Smith, 1967. Mate selection in a population of sockeye salmon (Oncorhynchus nerka) of mixed age groups. *J. Fish. Res. Bd. Canada* 24(9):1955-1977.
- Hayes, F.R., 1949. The growth, general chemistry and temperature relations of salmonid eggs. *Quart. Rev. Biol.* 24:281-300.
- Hayes, F.R., and F.H. Armstrong, 1942. Physical changes in the constituent parts of developing salmon eggs. *Can. J. Res., D*, 20:99-114.
- Hayes, F.R., and D. Pelluet, 1945. The effect of temperature on the growth and efficiency of yolk conversion in the salmo embryo. *Can. J. Res., D*, 23:7-15.
- Hayes, F.R., D. Pelluet, and E. Gorham, 1953. Some effects of temperature on the embryonic development of the salmon (Salmo salar). *Can. J. Zool.* 31:42-51.
- Hayes, F.R., I.R. Wilmot, and D.H. Livingstone, 1951. The oxygen consumption of the salmon egg in relation to development and activity. *J. Exp. Zool.* 116:377-395.
- Kinne, O., 1963. The effects of temperature and salinity on marine and brackish water animals. I. Temperature. *Oceanogr. Marine Biol. Ann. Rev.* 1:301-340.
- Kinne, O., and E.M. Kinne, 1962. Rates of development in a cyprinodont fish exposed to different temperature-salinity-oxygen combinations. *Can. J. Zool.* 40:231-253.